



Effects of Jasmonic acid on essential oil yield and chemical compositions of two Iranian landraces of basil (*Ocimum basilicum*) under reduced irrigation

Fatemeh Malekpoor^{1*}, Azam Salimi¹, Abdollah Ghasemi Pirbalouti^{2,3}

¹Department of Plant Biology, Faculty of Biology Science, Kharazmi University, 15719-14911, Tehran, Iran;

*Email: fmalekpoor87@yahoo.com or std_fmalekpoor@khu.ac.ir

²Medicinal Plants Department, Shahrekord Branch, Islamic Azad University, Shahrekord, 88146, Iran;

³Medicinal Plants Program, College of Natural Sciences, Massachusetts University, Amherst, 01003, MA, USA;

ARTICLE INFO

Type: Original Research

Topic: Physiology of Medicinal Plants

Received September 19th 2014

Accepted February 18th 2015

Key words:

- ✓ *Ocimum basilicum* L.
- ✓ Jasmonic acid
- ✓ Secondary metabolites
- ✓ Reduced irrigation

ABSTRACT

Background & Aim: Sweet basil (*Ocimum basilicum*), a plant that is extensively cultivated in some countries, is used to enhance the flavour of salads, sauces, pasta, confectioneries and other products as both a fresh and dried herb. The effect of foliar application of Jasmonic acid and reduced irrigation on essential oil yield and chemical components of two Iranian landrace of basil (*Ocimum basilicum*) were investigated.

Experimental: Treatments comprised control, 0.0, 200, and 400 μ L Jasmonic acid applied to plants under normal irrigation and stressed conditions (30 and 60% F.C) based on a completely randomized design with three replications. The essential oils from the aerial parts of basil were analyzed by GC-FID and GC/MS.

Results & Discussion: Results indicated that the different levels of Jasmonic acid and irrigation had significant effects on oil yield and some main components of the essential oil. The highest value of oil content was obtained from application of 400 μ L JA. Percentage of some chemical constituents in the essential oil extracted from the plants under stress was higher than non-stressed plants. Foliar application of Jasmonic acid significantly improved methyl chavicol in the oils, but reduced Germacrene- D, α -Cadinol and δ -Cadinene amounts.

Industrial and practical recommendations: Since, essential oil of basil, particularly methyl chavicol component, have many application in pharmaceutical and perfumery industry and has health benefits such as antiviral, antibacterial and antispasmodic activity, we can use Jasmonic acid elicitor to enhance the therapeutic properties of basil and also essential oil content and methyl chavicol of plant.

1. Introduction

Ocimum basilicum L. belongs to the family Lamiaceae is an herb that is extensively cultivated in some countries. Aerial parts, especially leaves of sweet basil

are widely used to enhance the flavour of foods such as salads, pasta, tomato products, vegetables, pizza, meat, soups, marine foods, confectioneries and other products (Ghasemi Pirbalouti, 2014). Traditionally basil has

been used as a medicinal plant in the treatment of headaches, coughs, diarrhea, constipation, warts, worms, and kidney malfunctions (Ghasemi Pirbalouti, 2014; Simon *et al.*, 1984). In Iranian traditional medicine, the aerial parts of the plant are perceived as a carminative, galactagogue, stomachic, and antispasmodic (Sajjadi, 2006). Moreover, basil possesses a range of biological properties, including action as an insect repellent and nematocide (Deshpande and Tipnis, 1997), having antibacterial (Hussain *et al.*, 2008), antiviral (Chiang *et al.*, 2005), antifungal activity (Hussain *et al.*, 2008), and antioxidant properties (Kwee and Niemeyer, 2011). Methyl chavicol, methyl cinnamate, methyl eugenol, citral, and linalool are generally the main chemotypes in sweet basil. Investigations (Sajjadi, 2006) on the chemical composition of the essential oil of basil, have demonstrated considerable variability.

In recent years, scientists have been looking for new alternatives to conventional methods for plant protection, and simultaneously for improvements in health properties and bioactive compound content. One of these methods is elicitation, which is induction of natural plant resistance mechanisms using biotic or abiotic factors. Elicitation can be an important strategy towards obtaining improved production of plant secondary metabolites *in vivo* (Hussain *et al.*, 2012). There are several commercially available chemical compounds that could be used as elicitors to modify plant secondary metabolites and subsequently the bioactivity of medicinal plants (Ghasemi Pirbalouti, *et al.*, 2014).

It has been reported that plant hormonal chemicals such as jasmonic acid (JA) and methyl jasmonate (MeJA), can be used as elicitors (Rabea *et al.*, 2003). Endogenous JA is a signaling compound that modulates various physiological processes in plants (Wasternack and Parthier, 1997). Therefore, exogenous JA has been used to induce useful secondary metabolites such as alkaloids, terpenoids, and phenolics in some plants such as *Nicotina* species, *Hyoscyamus muticus*, Norway spruce stems, and some plant cell cultures (Halitschke *et al.*, 2000; Keina *et al.*, 2001; Martin *et al.*, 2002). The increased amounts of secondary metabolites increase not only plants' disease resistance but also their market values regarding the increased health benefits. Moreover, environmental factors such as moisture deficit are known to play an

important role in the synthesis of secondary metabolites in plants (Ghasemi Pirbalouti *et al.*, 2015). A lot of researches (Khalid, 2006; Bettaieb, *et al.*, 2009; Ghasemi Pirbalouti *et al.*, 2013) showed the effect of water stress on secondary metabolites such as essential oils in different plant species. The present study was performed to evaluate the effect of various concentrations of Jasmonic acid on essential oil yield and essential oil composition two landraces of basil (*Ocimum basilicum* L.) under water deficit.

2. Materials and Methods

2.1. Plant material and experimental site description

Plastic pots were filled with clay loam at a pH of 7.23, containing 0.8% organic matter comprised of 0.01% total N, 11.20 mg/kg available phosphorus, 694 mg/kg available potassium, and a saline value measured at E.C.: 1.35 dS/m.

Two Iranian landraces of sweet basil (*Ocimum basilicum* L.) seeds were obtained from the Pakan Seed Company, Isfahan, Iran. In May 2014, pots were transferred to the field in Shahrekord (latitude 32° 20' N, longitude 50° 51' E, altitude 2061 m above sea level), southwestern Iran.

Type of study area climate by Emberger's climatology method is cold and semiarid and semi humid with temperate summer and very cold winter by Karimi's climatology method (IRIMO, 2012). In this study, no inorganic fertilizer and systemic pesticide was used during the entire experiment, and weed control was done manually.

2.2. Experimental design and treatments

Treatments included three irrigation regimes, viz., I₁ (unstressed or control), I₂ (irrigation in 70% field capacity when 30% of maximum total available soil water was depleted in the upper 30 cm of the soil profile), I₃ (irrigation in 40% field capacity when 60% of maximum total available soil water was depleted in the upper 30 cm of the soil profile) and three Jasmonic acid treatments (0.0, 200 and 400 µl) sprayed thrice at 10-12 leaves, before flowering, and two weeks later. Jasmonic acid was dissolved in ethanol 80%, diluted in distilled water with various concentrations. These solutions were sprayed at dew point (approximately 100 ml per plant) with a hand sprayer (untreated control plants were sprayed with an equivalent volume

of distilled water). The aerial parts of basil were collected after third spray of Jasmonic acid in July 2014.

2.3. Essential oil isolation

Dried plant material (100 g) was powdered and subjected to hydro-distillation for three hours using a Clevenger-type apparatus ([British Pharmacopoeia, 1988](#)). The essential oils were dried with anhydrous sodium sulphate and kept in amber vials at 4 °C prior to use.

2.4. Identification of the oil constituents

Composition of the essential oils was determined by gas chromatography (GC) and mass spectrophotometry (GC/MS). The GC analysis was done on an Agilent Technologies 7890 GC (Agilent Technologies, Santa Clara, CA) equipped with a single injector and a flame ionization detector (FID). A polar HP Innowax column and an apolar HP-5 capillary column (30 m x 0.25 mm, 0.25 µm film thicknesses) coated with 5% phenyl, 95% methyl polysiloxane were used. The flow of the carrier gas (N₂) was 0.8 ml/min. Initial column temperature was 60°C and programmed to increase at 4°C/min to 280°C. The injector temperature was set at 280 and 300°C. Split injection was conducted with a ratio split of 1:40. Essential oil samples of 0.1 µL were injected neat (directly).

GC-MS analyses of aromatic oil samples were performed on an Agilent Technologies 7890 gas chromatograph coupled to Agilent 5975 C mass selective detector (MSD) and quadrupole EI mass analyzer (Agilent Technologies, Palo Alto, CA, USA). A HP-5MS 5% column (coated with methyl silicone) (30 m x 0.25 mm, 0.25 µm film thicknesses) was used as the stationary phase. Helium was used as the carrier gas at 0.8 mL/min flow rate. The temperature was programmed from 60 to 280°C at 4°C/min ramp rate. The injector and the GC-MS interface temperatures were maintained at 290°C and 300°C, respectively. Mass spectra were recorded at 70 eV. Mass range was from *m/z* 50-550. The ion source and the detector temperatures were maintained at 250 and 150°C, respectively.

Oil constituents were identified based on their retention indices (determined with reference to homologous series of C₅-C₂₄ *n*-alkanes), by comparison of their mass spectra with those reported in the literature ([Adams, 2007](#)) and stored in NIST 08

(National Institute of Standards and Technology) and Willey (ChemStation data system) libraries. The peak area percentages were computed from HP-5 column without the use of FID response factors.

2.5. Statistical analysis

Simple and interaction effects of experimental factors were derived from two-way analysis of variance (ANOVA) based on the GLM procedure of the SAS statistical package (SAS/STAT® v.9.2. SAS Institute Inc., Cary, NC).

The assumptions of variance analysis were tested by ensuring that the residuals were random and homogenous, with a normal distribution about a zero mean. The significance of differences among treatment means was tested using Duncan's multiple range test (DMRT) at $p \leq 0.05$.

3. Results and discussion

3.1. Essential oil yield

All essential oils extracted from the aerial parts of two landraces of basil under different treatments were clear and yellow liquid. An analysis of variance indicated that different irrigation regimes had a significant effect on oil yield of basil ($p \leq 0.01$) ([Table 1](#)). The highest oil yield was obtained from mild drought stress treatment (I₃) with 0.82 ml/100 g dry material ([Table 1](#)). Our results indicated that Jasmonic acid at 400 µl produced maximum oil content (0.53 ml/100 g) ([Table 1](#)). It has been demonstrated that signal molecules such as Jasmonic acid are very potential elicitors for induction of plant secondary metabolites ([Zhao *et al.*, 2005](#)). Recent years, the applications of signal components as elicitors have evolved an effective strategy for the production of target secondary metabolites in plant. Foliar application of Jasmonic acid on oil content of basil was influenced ($p \leq 0.01$) by irrigation regime ([Table 2](#)). The highest oil yield (0.89 ml/100 g) was achieved by foliar spray of 400 µL Jasmonic acid under reduced irrigation or mild stress drought (I₃ × J₄) ([Table 2](#)). The lowest oil content was obtained when Jasmonic acid was applied under normal irrigation ([Table 2](#)).

There was significant jasmonic acid × landrace interaction effect on oil content ([Table 3](#)). In this study, landrace × irrigation had significant effects on oil content ($p \leq 0.01$) ([Table 4](#)).

Table 1. The effects of irrigation regime, jasmonic acid and landrace on the essential oil from basil

Treatments	Oil yield (ml/100 g)	1,8-Cineole	Linalool	Methyl chavicol	Nerol	Neral	Geranial	β -Caryophyllene	α -Humulene	Germacrene-D	δ -Cadinene	Caryophyllene oxide	α -Cadinol
Irrigation													
I1 (control)	0.183c	1.120b	4.39	33.24a	1.44b	6.30	7.10	4.82	3.12ab	2.98a	1.79a	1.854b	5.92a
I2 (60% FC)	0.488b	1.349a	4.41	34.71ab	1.77a	5.93	6.93	4.55	2.88b	2.66b	1.59b	1.79b	4.79b
I3 (30% FC)	0.822a	1.478a	4.62	35.67a	1.51b	5.74	6.82	4.99	3.16a	2.43b	1.41c	2.24a	5.42a
ANOVA	$p \leq 0.01$	$p \leq 0.01$	ns	$p \leq 0.05$	$p \leq 0.01$	ns	ns	ns	$p \leq 0.05$	$p \leq 0.01$	$p \leq 0.01$	$p \leq 0.01$	$p \leq 0.01$
Jasmonic acid													
J1 (control)	0.471c	1.317a	4.44a	34.55ab	1.63ab	6.39a	7.36a	0.49ab	0.99a	4.98a	3.35a	2.70ab	1.68ab
J2 (ethanol)	0.467d	1.316a	4.23a	32.91b	1.51b	5.81ab	6.65a	0.50a	1.09a	5.15a	3.18ab	2.89a	1.78a
J3 (200)	0.518b	1.251a	4.48a	34.41ab	1.79a	6.26ab	7.27a	0.44b	0.99a	4.88a	2.98bc	2.66ab	1.54bc
J4 (400)	0.535a	1.378a	4.74a	36.29a	1.28c	5.50b	6.51a	0.45ab	0.98a	4.14b	2.71c	2.51b	1.39c
ANOVA	$P \leq 0.01$	ns	ns	$P \leq 0.05$	$P \leq 0.01$	ns	ns	ns	ns	$P \leq 0.01$	$P \leq 0.01$	ns	$P \leq 0.01$
Landraces													
S1 (Green)	0.71a	0.321b	1.50b	28.10b	2.84a	10.83a	12.64a	6.86a	3.79a	2.65	0.32b	2.85a	2.12b
S2 (Purple)	0.285b	2.31a	7.45a	40.98a	0.27b	1.14b	1.25b	2.71b	2.33b	2.73	2.87a	1.06b	8.64a
ANOVA	$p \leq 0.01$	$p \leq 0.01$	$p \leq 0.01$	$p \leq 0.01$	$p \leq 0.01$	$p \leq 0.01$	$p \leq 0.01$	$p \leq 0.01$	$p \leq 0.01$	ns	$p \leq 0.01$	$p \leq 0.01$	$p \leq 0.01$

ns: non-significant

Table 2. The interaction effects of irrigation regime \times jasmonic acid on the essential oil from basil

Interaction effect	Oil yield	1,8-Cineole	Linalool	Methyl chavicol	Nerol	Neral	Geranial	β -Caryophyllene	α -Humulene	Germacrene-D	δ -Cadinene	Caryophyllene oxide	α -Cadinol
I1 \times J1	0.175i	1.00cd	3.84c	29.34e	1.46cd	6.28ab	6.88ab	5.56a	3.52a	3.47a	2.31a	2.15ab	7.28a
I1 \times J2	0.175i	1.22bc	4.72abc	34.63bcd	1.42cde	6.33ab	7.11ab	5.03ab	3.30abc	2.86bc	1.76bc	1.94a-d	6.02b
I1 \times J3	0.179i	0.80d	3.94bc	35.21bcd	1.44cde	6.35ab	7.24ab	4.77abc	3.05abcde	2.96b	1.52cdef	1.84bcde	4.94bcd
I1 \times J4	0.204h	1.44ab	5.06a	33.79bcd	1.44cd	6.23ab	7.16ab	3.94cd	2.61ef	2.63bcd	1.58cde	1.43e	5.44bc
I2 \times J1	0.450g	1.42ab	4.36abc	32.09de	1.70abc	5.64ab	6.59ab	4.62bcd	2.80-f	2.70bcd	1.60cd	1.56cde	4.81cd
I2 \times J2	0.462f	1.19bc	3.88bc	32.12de	1.87ab	6.76a	7.90a	4.82abc	3.35abc	2.88bc	1.76bc	2.04abc	4.95bcd
I2 \times J3	0.525d	1.33abc	4.69abc	34.85bcd	1.95ab	6.09ab	7.24ab	4.91ab	2.89c-f	2.55bcd	1.53cdef	2.08ab	5.31bc
I2 \times J4	0.513e	1.44ab	4.70abc	39.80a	1.32de	5.21b	5.99b	3.85d	2.51f	2.52bcd	1.32ef	1.50de	4.08d
I3 \times J1	0.775c	1.51ab	4.49abc	37.30ab	1.36cde	5.50ab	6.47ab	5.28ab	3.22a-d	2.51bcd	1.43def	2.40a	5.56bc
I3 \times J2	0.775c	1.53ab	4.73abc	36.91abc	1.61bcd	6.07ab	7.07ab	5.08ab	3.42ab	2.37d	1.38def	2.26a	5.37bc
I3 \times J3	0.850b	1.60a	4.81ab	33.15cde	1.99a	6.32ab	7.32ab	4.96ab	3.01b-e	2.47cd	1.57cde	1.96a-d	5.46bc
I3 \times J4	0.888a	1.25bc	4.45abc	35.29bcd	1.08e	5.05b	6.39ab	4.65bcd	3.01b-e	2.38d	1.26f	2.36a	5.31bc
ANOVA	$p \leq 0.01$	$p \leq 0.01$	ns	$p \leq 0.01$	$p \leq 0.05$	ns	ns	ns	ns	ns	$p \leq 0.05$	$p \leq 0.05$	$p \leq 0.05$

Table 3. The interaction effects of jasmonic acid × landrace on the essential oil from basil.

Interaction effect	Oil yield	1,8-Cineole	Linalool	Methyl chavicol	Nerol	Neral	Geranial	β-Caryophyllene	α-Humulene	Germacrene - D	δ-Cadinene	Caryophyllene oxide	α-Cadinol
J1 × S1	0.683c	0.25b	1.20c	28.92d	2.74b	10.46a	12.07a	7.33a	3.96a	2.81ab	0.33d	2.94a	2.22c
J1 × S2	0.250g	2.38a	7.26a	36.90c	0.27d	1.15b	1.23b	2.98c	2.40d	2.98a	3.24a	1.14bc	9.54a
J2 × S1	0.683c	0.24b	1.24c	28.37d	2.94b	11.46a	13.26a	7.08a	3.82ab	2.79ab	0.31d	2.80a	2.12c
J2 × S2	0.258f	2.39a	7.65a	40.73b	0.32d	1.31b	1.46b	2.87cd	2.89c	2.62ab	3.05a	1.36b	8.77ab
J3 × S1	0.733b	0.31b	1.55bc	28.95d	3.26a	10.99a	12.87a	6.93a	3.80ab	2.50b	0.32d	2.93a	2.11c
J3 × S2	0.303e	2.19a	7.40a	39.86bc	0.33d	1.52b	1.67b	2.83cd	2.17de	2.81ab	2.75b	0.99bc	8.36b
J4 × S1	0.742a	0.47b	2.00b	26.15d	2.42c	10.41a	12.37a	6.11b	3.56b	2.51b	0.31d	2.74a	2.01c
J4 × S2	0.328d	2.27a	7.47a	46.43a	0.14d	0.59b	0.65b	2.17d	1.85e	2.50b	2.46c	0.78c	7.87b
ANOVA	<i>p</i> ≤ 0.01	<i>ns</i>	<i>ns</i>	<i>p</i> ≤ 0.01	<i>p</i> ≤ 0.05	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>p</i> ≤ 0.05	<i>ns</i>	<i>p</i> ≤ 0.01	<i>ns</i>	<i>ns</i>

Table 4. The interaction effects of irrigation regime × landrace on the essential oil from basil.

Interaction effect	Oil yield	1,8-Cineole	Linalool	Methyl chavicol	Nerol	Neral	Geranial	β-Caryophyllene	α-Humulene	Germacrene-D	δ-Cadinene	Caryophyllene oxide	α-Cadinol
I1 × S1	0.225d	0.37d	1.72c	27.71cd	2.72b	11.35a	12.98a	6.65b	3.71b	2.75b	0.31d	2.67b	2.22cd
I1 × S2	0.142e	1.86c	7.05b	38.77b	0.16c	1.25b	1.22b	3.00c	2.54c	3.21a	3.28a	1.01c	9.62a
I2 × S1	0.750b	0.29d	1.52c	30.37c	3.06a	10.49a	12.22a	6.33b	3.45b	2.73b	0.35d	2.37b	1.54d
I2 × S2	0.225d	2.40b	7.29b	39.05b	0.36c	1.36e	1.63b	2.77c	2.32cd	2.60bc	2.83b	1.22c	8.04b
I3 × S1	1.15a	0.29d	1.25c	26.21d	2.74b	10.65a	12.72a	7.61a	4.20a	2.48bc	0.30d	3.52a	2.59c
I3 × S2	0.488c	2.66a	1.99a	45.12a	0.28c	0.82b	0.91b	2.37c	2.13d	2.38c	2.52c	0.96c	8.26b
ANOVA	<i>p</i> ≤ 0.01	<i>p</i> ≤ 0.01	<i>p</i> ≤ 0.01	<i>p</i> ≤ 0.01	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>p</i> ≤ 0.01	<i>p</i> ≤ 0.01	<i>p</i> ≤ 0.05	<i>p</i> ≤ 0.01	<i>p</i> ≤ 0.01	<i>p</i> ≤ 0.05

Table 5. The interaction effects of irrigation regime \times jasmonic acid \times landrace on the essential oil from basil.

Interaction effect	Oil yield	1,8-Cineole	Linalool	Methyl chavicol	Nerol	Neral	Geranial	β -Caryophyllene	α -Humulene	Germacrene-D	δ -Cadinene	Caryophyllene oxide	α -Cadinol
I1 \times J1 \times S1	0.225k	0.34h	1.32fg	23.70jk	2.74cde	11.28abc	12.65a-d	7.71ab	4.13ab	3.19abc	0.33g	2.98bcd	1.92fgh
I1 \times J1 \times S2	0.125n	1.67fg	6.35de	34.99de	0.17gh	1.28de	1.12e	3.40gh	2.91efg	3.75a	4.30a	1.33gh	12.63a
I1 \times J2 \times S1	0.225k	0.40h	1.63fg	27.23g-j	2.64cde	11.48ab	13.07a-d	7.07a-d	3.85a-d	2.89b-f	0.31g	2.79cde	2.12gh
I1 \times J2 \times S2	0.125n	2.04ef	7.80abc	42.03bc	0.20gh	1.18de	1.15e	3.00g-j	2.77fgh	2.83b-f	3.21bc	1.09hi	9.93b
I1 \times J3 \times S1	0.0200l	0.21h	1.95fg	33.15def	2.60de	10.34abc	12.04a-d	5.78ef	3.42c-f	2.61c-f	0.32g	2.57c-f	2.87g
I1 \times J3 \times S2	0.158m	1.39g	5.92e	37.28cd	0.27gh	2.37d	2.44e	3.76g	2.69ghi	3.31ab	2.72de	1.11hi	7.01e
I1 \times J4 \times S1	0.250j	0.52h	1.98fg	26.78h-k	2.88bcd	12.32a	14.16ab	6.05def	3.44c-f	2.32def	0.29g	2.33def	1.98fgh
I1 \times J4 \times S2	0.158m	2.35cde	8.13ab	40.79bc	0.01h	0.15e	0.16e	1.82j	1.77k	2.95b-e	2.88cd	0.53i	8.89bcd
I2 \times J1 \times S1	0.700f	0.15h	1.11fg	32.80de	3.13bc	10.26abc	11.91a-d	6.44c-f	3.42c-f	2.81b-f	0.33g	2.08f	1.73fgh
I2 \times J1 \times S2	0.200l	2.68a-d	7.62a-d	31.38e-h	0.28gh	10.03de	1.26e	2.79g-j	2.17h-k	2.60c-f	2.88cd	1.04hi	7.89cde
I2 \times J2 \times S1	0.700f	0.12h	0.99g	29.51e-i	3.31ab	11.73ab	13.64abc	6.52b-f	3.46b-e	2.96bcd	0.31g	2.10ef	1.56gh
I2 \times J2 \times S2	0.225g	2.26de	6.77cde	34.72de	0.43gh	1.78de	2.15e	3.11ghi	3.23d-g	2.80b-f	3.49b	1.98fg	8.33b-e
I2 \times J3 \times S1	0.850d	0.42h	1.58ab	38.83f-j	3.34ab	10.25abc	12.21a-d	7.04a-e	3.71a-d	2.39def	0.37g	3.12abc	2.09fgh
I2 \times J3 \times S2	0.200l	2.25de	7.79abc	40.88bc	0.56g	1.93de	2.27e	2.77g-j	2.07ijk	2.71b-f	2.69de	1.03hi	8.53b-e
I2 \times J4 \times S1	0.750e	0.47h	2.41f	30.36e-i	2.47de	9.71bc	11.13d	5.30f	3.22d-g	2.76b	0.38g	2.18ef	0.77h
I2 \times J4 \times S2	0.275i	2.41b-e	6.99b-e	49.23a	0.16gh	0.71de	0.85e	2.40hij	1.80k	2.28ef	2.25f	0.81hi	7.39de
I3 \times J1 \times S1	1.125c	0.24h	1.16fg	30.27e-i	2.36ef	9.85bc	11.64cd	7.83a	4.32a	2.42def	0.33g	3.76a	3.02fg
I3 \times J1 \times S2	0.425h	2.79abc	7.82abc	44.34ab	0.36gh	1.15de	1.31e	2.73g-j	2.12h-k	2.59c-f	2.53def	1.04hi	8.09cde
I3 \times J2 \times S1	1.125c	0.20h	1.10g	28.38f-j	2.87bcd	11.17abc	13.06a-d	7.65abc	4.17a	2.51c-f	0.31g	3.52ab	2.67fg
I3 \times J2 \times S2	0.425h	2.86ab	8.36a	45.44ab	0.35gh	0.97de	1.09e	2.50g-j	2.66g-j	2.23f	2.45ef	0.99hi	8.06cde
I3 \times J3 \times S1	1.150b	0.29h	1.12fg	24.87ijk	3.82a	12.38a	14.35a	7.96a	4.28a	2.52cdef	0.28g	3.09abc	1.37gh
I3 \times J3 \times S2	0.550g	2.92a	8.49a	41.44bc	0.17gh	0.26de	0.30e	1.95ij	1.74k	2.42def	2.86cd	0.82hi	9.54bc
I3 \times J4 \times S1	1.225a	0.43h	1.61fg	21.32k	1.92f	9.19c	11.84bcd	6.99a-e	4.03abc	2.46def	0.27g	3.72a	3.28f
I3 \times J4 \times S2	0.550g	2.06ef	7.29a-d	49.27a	0.24gh	0.91de	0.94e	2.30hij	1.99jk	2.29def	2.24f	0.99hi	7.34de
ANOVA	$p \leq 0.01$	$p \leq 0.05$	$p \leq 0.05$	$p \leq 0.01$	$p \leq 0.01$	$p \leq 0.05$	$p \leq 0.01$	<i>ns</i>	<i>ns</i>	<i>ns</i>	$p \leq 0.01$	$p \leq 0.05$	$p \leq 0.01$

The highest oil yield was observed in the green landrace of basil under reduced irrigation or mild stress drought (Table 4). In this study, jasmonic acid \times irrigation \times landrace had significant effects ($p \leq 0.01$) on oil yield (Table 5). The highest oil yield (1.22 ml / 100 g dry weight) was achieved by foliar spray of jasmonic at 400 μ l under reduced irrigation in the green landrace of basil (Table 5).

Similarly, Simon *et al.* (1992) reported that water stress increased essential oil accumulation via higher density of oil glands due to the reduction in leaf area. Moreover, Shokrani *et al.* (2012) reported an increase in essential oil content and yield in basil with increasing water stress severity and higher yield of essential oil obtained from mild water stress. In general, moisture deficit increases the essential oil content of more medicinal and aromatic plants, because in case of stress, more metabolites are produced in the plants and substances prevent from oxidization in the cells. Essential oil helps to plant that easier adopt to the environmental stress conditions. In addition, stress induced alterations in oil accumulation are considered to be mainly due to its effect on plant growth and differentiation, due to the fact that plants produce high terpene concentrations under water deficit conditions due to a low allocation of carbon to the growth, suggesting a trade-off between growth and defense (Turtola *et al.*, 2003). Probably, improvement in oil yield by exogenous application of Jasmonic acid might be due to the increase in cycle growth, nutrients uptake or changes in leaf oil gland population and monoterpenes biosynthesis (Ghasemi Pirbalouti *et al.*, 2014). Boonlertnirun *et al.* (2008) found that foliar application of elicitor had a significant effect on the growth, and yield of drought-stressed rice plants compared to control plants.

3.2. Chemical compositions of oil

The chemical constituents of the essential oil identified by GC-FID and GC/MS. Tables 1-5 show the effect of different treatments on the chemical composition of essential oil extracted from two landraces of basil. The major components in the essential oil were methyl chavicol, geraniol, nerol, and β -caryophyllene (Table 1). Our results demonstrated that the essential oils obtained from the landraces of basil contained phenylpropanoids (methyl chavicol), oxygenated monoterpenes (linalool, geraniol, and nerol),

sesquiterpenes hydrocarbons (β -caryophyllene), and oxygenated sesquiterpenes (α -cadinol), confirming earlier reports that major chemical groups obtained from the aerial parts of basil were phenylpropanoids and oxygenated monoterpenes (Sajjadi, 2006; Ghasemi Pirbalouti *et al.*, 2013; Ghasemi Pirbalouti, 2014).

An analysis of variance indicated that the different factors had a significant effect on some major constituents of the essential oil from basil. Results also indicated that there were significant differences between Jasmonic acid, and landraces treatments in term methyl chavicol (Table 1). Results of mean comparison revealed that amount of methyl chavicol, as a main compound in the basil oil, not considerably affected by drought levels (Table 1). Different levels of the foliar application of Jasmonic acid had significant effects on the major constituents of the essential oils (Table 1). This demonstrated that Jasmonic acid at 400 μ L produced the highest levels of methyl chavicol (36.29%), linalool (4.74%), and 1,8-Cineole (1.37%), but at 200 μ L produced the highest levels of nerol (1.79%), geraniol (0.94%) (Table 1).

Results of interaction effects of irrigation \times Jasmonic acid indicated that the highest contents of some main components were obtained from the application of Jasmonic acid under reduced irrigation (Table 2). Our results further indicated that foliar application of Jasmonic acid reduced the effect of water deficit stress on amounts of linalool, methyl chavicol, and geraniol, the major compounds in the essential oil from the landraces of basil (Table 2). Jasmonic acid may play an important signaling role in the activation of various plant defense responses, such as the biosynthesis of special secondary metabolites.

In present research, the ANOVA showed that interaction effects of landrace \times jasmonic acid had significant effects on the major constituents of basil essential oils (Table 3). In addition, interaction effects of irrigation \times landrace had significant effects on many volatile constituents (Table 4). Moreover, interaction effects of jasmonic acid \times irrigation \times landrace had significant effects on many volatile constituents (Table 5). The maximum amount of methyl chavicol (49.27%) was obtained by 400 μ l jasmonic acid in the purple landrace under mild drought stress.

4. Conclusions

In this research, effects of foliar application of jasmonic acid and drought stress on yield and chemical composition of essential oils of two landraces of basil have evaluated. The investigation showed two landraces of basil are sensitive to the application of elicitors such as jasmonic acid and water deficit. The results revealed that essential oil yield two landraces of basil increased with application of elicitors under drought stress. Not only oil ratio but also oil composition was affected since some components were increased and other components were decreased. Finally, it could be concluded that foliar application of jasmonic acid and deficit irrigation, as a possible technique, can be used to increase volatile oil ratio in basil cultivation. This study provides the basis for further research on improving the quality and pro-health functional value of two landraces of basil using elicitors.

5. References

- Adams, R.P. 2007. Identification of essential oil components by gas chromatography/quadrupole mass spectroscopy. Allured Publishing Corporation, CarolStream, IL, USA.
- Bettaieb, I., Zakhama, N., Wannes, W.A., Kchouk, M. E., Marzouk, B. 2009. Water deficit effects on *Salvia officinalis* fatty acids and essential oils composition. *Scientia Horticulturae*, 120: 271–275.
- Boonlertnirun, S., Boonraung, C., Suvanasa, R. 2008. Application of chitosan in Rice production. *International Journal of Minerals, Metallurgy, and Materials*, 18: 47–52.
- British Pharmacopoeia, 1988. British Pharmacopoeia, vol. 2. HMSO, London, pp.137–138.
- Chiang, L.C., Ng, L.T., Cheng, P.W., Chiang, W., Lin, C.C. 2005. Antiviral activities of extracts and selected pure constituents of *Ocimum basilicum*. *Clinical and Experimental Pharmacology and Physiology*, 32(10): 811–816.
- Deshpande, R.S., Tipnis, H.P. 1997. Insecticidal activity of *Ocimum basilicum* L. *Pesticides*, 11, 1–12.
- Ghasemi Pirbalouti, A., Samani, M.R., Hashemi, M., Zeinali, H. 2013. Salicylic acid affects growth, essential oil and chemical compositions of thyme (*Thymus daenensis* Celak.) under reduced irrigation. *Plant Growth Regulation*, 72: 289–301.
- Ghasemi Pirbalouti, A. 2014. Diversity in chemical composition and yield of essential oil from two Iranian landraces of sweet basil. *Genetika*, 46: 419–426.
- Ghasemi Pirbalouti, A., Sajjadi, S.E., Parang, K. 2014. A review (research and patents) on jasmonic acid and its derivatives. *Archiv der Pharmazie*, 347: 229–239.
- Ghasemi Pirbalouti, A., Ghahfarokhi, B., Ghahfarokhi, S.A.M., Malekpoor, F. 2015. Chemical composition of essential oils from the aerial parts and underground parts of Iranian valerian collected from different natural habitats. *Industrial Crops and Products*, 63: 147–151.
- Halitschke, R., Kessler, A., Kahl, J., Lornez, A., Baldwin, I.T. 2000. Ecophysiological comparison of direct and indirect defenses in *Nicotiana attenuata*. *Oecologia*, 124: 408–417.
- Hussain, A.I., Anwar, F., Hussain Sherazi, S.T., Przybylski, R. 2008. Chemical composition, antioxidant and antimicrobial activities of basil (*Ocimum basilicum*) essential oils depends on seasonal variations. *Food Chemistry*, 108(3): 986–995.
- Hussain, S., Fareed, S., Ansari, S., Rahman, A., Ahmad, I.Z., Saeed, M. 2012. Current approaches toward production of secondary plant metabolites. *Journal of Pharmacy and Bioallied Sciences*, 4: 10–20.
- IRIMO, 2012. (Islamic Republic of Iran Meteorological Organization). Climatology methods. <http://www.irimo.ir>.
- Keina, M., Oldham, N., Baldwin, I.T. 2001. Rapid HPLC screening of jasmonate-induced increases in tobacco alkaloids, phenolics, and diterpene glycosides in *Nicotiana attenuate*. *Journal of Agricultural and Food Chemistry*, 49: 3553–3558.
- Khalid, K.A. 2006. Influence of water stress on growth, essential oil and chemical composition of herbs (*Ocimum sp.*). *International Agrophysics*, 20: 289–296.
- Kwee, E.M., Niemeyer, E.D. 2011. Variations in phenolic composition and antioxidant properties among 15 basil (*Ocimum basilicum* L.) cultivars. *Food Chemistry*, 128(4): 1044–1050.

- Martin, D., Tholl, D., Gershenzon, J., Bohlmann, J. 2002. Methyl jasmonate induces traumatic resin ducts, terpenoid resin biosynthesis, and terpenoid accumulation in developing xylem of Norway spruce stems. *Plant Physiology*, 129: 1003-1018.
- Rabea, E.I., Badawy, M.E., Stevens, C., Smaghe, G., Steurbaut, W. 2003. Chitosan as antimicrobial agent: applications and mode of action. *Biomacromolecules*, 4: 1457-1465.
- Sajjadi, S.E. 2006. Analysis of the essential oils of two cultivated basil (*Ocimum basilicum* L.) From Iran. *DARU Journal of Pharmaceutical Sciences*, 14: 128-130.
- Shokrani, F., Pirzad, A., Zardoshti, M., Darvishzadeh, R. 2012. Effect of biological nitrogen on the yield of dried flower and essential oil of *Calendula officinalis* L. under end season water deficit condition. *International Research Journal of Applied and Basic Sciences*, 3: 24-34.
- Simon, J.E., Reiss-Bubenheim, D., Joly, R.J., Charles, D.J. 1992. Water stress-induced alterations in essential oil content and composition of sweet basil. *Journal of Essential Oil Research*, 4: 71-75.
- Simon, J.E., Chadwick, A.F., Craker, L.E. 1984. Herbs: an indexed bibliography, 1971-1980. The scientific literature on selected herbs, and aromatic and medicinal plants of the temperate zone. Archon Books: Hamden, CT.
- Turtola, S., Manninen, A.M., Rikala, R., Kainulainen, P. 2003. Drought stress alters the concentration of wood terpenoids in Scots pine and Norway spruce seedlings. *Journal of Chemical Ecology*, 29: 1981-1985.
- Wasternack, C., Parthier, B. 1997. Jasmonate-signaled plant gene expression. *Trends Plant Science*, 2: 1360-1385.
- Zhao, J., Davis, L.C., Verpoorte, R. 2005. Elicitor signal transduction leading to production of plant secondary metabolites. *Biotechnology Advances*, 23: 283-333.

