



Phytochemical investigation of the essential oils from the leaf and stem of *Dorema ammoniacum* D. Don. (Apiaceae) in Central Zagros, Iran

Fakhteh Zandpour¹, Mohammad Reza Vahabi^{1*}, Ali Reza Allafchian², Hamid Reza Farhang³

¹Department of Natural Resources, Isfahan University of Technology, Isfahan 84156–83111, Iran;

*Email: Vahabi@cc.iut.ac.ir

²Research Institute for Nanotechnology and Advanced Materials, Isfahan University of Technology, Isfahan 84156–83111, Iran;

³MSc of Range Management, Isfahan University of Technology, Isfahan 84156–83111, Iran;

ARTICLE INFO

Type: Original Research

Topic: Medicinal Plants

Received February 19th 2016

Accepted April 2th 2016

Key words:

- ✓ *Dorema ammoniacum* D. Don.
- ✓ Essential oil
- ✓ GC/MS
- ✓ Isfahan province
- ✓ Iran

ABSTRACT

Background & Aim: *Dorema ammoniacum* D. Don. (ammoniac plant) belongs to Apiaceae family and *Dorema* genus. It is one of the most important endemic medicinal plants in its family which grows in the arid and semi-arid areas in the regions of Iran and some of the Asian countries such as Turkmenistan, Afghanistan, Pakistan and India. This study was carried out with aim of the extraction and identification of constituents of *D. ammoniacum* D. Don. in the western areas of Isfahan province, Iran.

Experimental: For this purpose, after the plant collecting in the mentioned area its target organs namely stem and leaf were separated, air dried, milled and weighted in the certain amount. Then the essential oil of the plant was isolated by hydrodistillation method and analyzed by GC/MS device.

Results: The results indicated that major components were Limonene (49.18%), Neophytadiene (10.21%), beta Caryophyllene (3.54%), Phytol (2.61%) and Neryl acetone (1.46%).

Recommended applications/industries: Several therapeutic cases have been reported in traditional medicine and currently in contemporary medicine for *Dorema ammoniacum* D. Don. that some of them are include the treatment of cough, vasodilator, anti-flatulence and anti-microbial. Also, the mentioned plant has varied applications in the fields of foods, cosmetic and detergent industries.

1. Introduction

Essential oils have been used for thousands of years in various cultures for medicinal and health purposes. They have a widespread range of application from aromatherapy, household cleaning products, personal beauty care and natural medicine treatments (Worwood, 1991; Farhang et al., 2015). The Apiaceae family has

been distributed with 300 genera and 3000 plant species in the world approximately. The genus *Dorema* D. Don (Apiaceae) is represented in the flora of Iran by seven species, among which both *D. ammoniacum* D. Don. and *D. aucheri* Boiss. are endemic (Mozaffarian, 1996). *Dorema ammoniacum* D. Don. (ammoniac plant) is a perennial monocarpic plant grows to height about 1–3 m and in spring and early summer contains a milky juice. It is one of the most important endemic medicinal plants

in its family which grows in the arid and semi-arid areas in the regions of Iran and some of the Asian countries such as Turkmenistan, Afghanistan, Pakistan and India (Rechinger, 1980). The mentioned species in the Iranian traditional medicinal has numerous names. Some of the most prevalent names include ushaq, vasha, kandal, koma-kandal and also as Persian ammoniacum in Greek and medicinal literatures (Amin, 1991; Mozaffarian, 1996; Rechinger, 1980; Howes, 1950). *Dorema ammoniacum* D. Don. produces a medicinal gum resin, commonly known as ammoniacum gum, which can be found in the holes of stems, roots, and petioles (Langenheim, 2003). Overall, the plants extract constituents and their secondary metabolisms are provided use to efficacious on human and animal health that can be used to cure maladies (Evans, 1996). The gum resin of *Dorema ammoniacum* D. Don. in the Iranian traditional medicine has been considered beneficial in treatment of spastic pains, gastric disorders, intestinal parasitic infections and skin inflammations and as analgesic, stimulant, expectorant and laxative (Amin, 1991; Parsa, 1959; Zarshenas *et al.*, 2013). In addition the existence resin in the stem of *Dorema ammoniacum* D. Don. are used as antimicrobial, vasodilator, a carminative, diaphoretic, mildly diuretic, expectorant and poultice (Irvani *et al.*, 2010). *Dorema ammoniacum* D. Don. in the British pharmacopeia as the plant with various therapeutic properties has been recognized that some of the most important of them are as follow: antispasmodic, sputum containing horny sputum, cough irritant, sudorific, the regulation of menstrual, reducer of blood fats, prevention of diabetes, hepatoprotective mechanism, sexual stimulus and sedative (British herbal pharmacopoeia, 1983; British pharmacopoeia, 1993). Several studies have been conducted in relation to identification of chemical compounds of the gum and volatile oil of different parts of *Dorema ammoniacum* D. Don. in some mentioned countries (Yassa *et al.*, 2014; Yousefzadi *et al.*, 2011a; Yousefzadi *et al.*, 2011b; Sajjadi *et al.*, 2007; Appendino *et al.*, 1991; and Arnone *et al.*, 1991). In all these studies miscellaneous chemical compositions were identified and the amount of the yields of essential oils was varied. The mentioned plant species in France are utilized in the factories producing of glues and cosmetic industry. Moreover, this plant has other varied uses such as food industry, confectionary products, paint industry, manufacturing, and production of the detergents and

especially in production of special glues for jewelry (Bown, 1995; Grieve, 1971; Chevallier, 2001). Plants, with their wide variety of chemical constituents, offer a promising source of new antimicrobial agents with general as well as specific antimicrobial activity (Evans, 1996). In the previous studies the antimicrobial activity of the extract of *Dorema ammoniacum* D. Don. had been reported (Rajani, 2002). The purpose of the current study was to identify all chemical compositions of the essential oil of *Dorema ammoniacum* D. Don. in the western areas of Isfahan province which is situated in the section of the Central Zagros area, Iran; for the first time.

2. Materials and Methods

2.1. Plant materials

Dorema ammoniacum D. Don. (Apiaceae) is perennial endemic native herb plant that grows as high as three meters (around one to three meters) in Iran, between the months of March and June. Samples of *Dorema ammoniacum* D. Don. aerial parts consisted of the leaves and stems collected in the middle of March 2015 (in its natural habitat situated in the Bardasiab zone of the Fereyduh Shahr county which is part of Central Zagros area, Iran), during the flowering period. The taxonomic identity of the plant was confirmed by comparing the collected voucher specimen with that of the known identity available in the herbarium of the Department of Natural Resources, Isfahan University of Technology (IUT), Iran (Fig. 1).

2.2. Experimental procedure

At first, 1000g of the aerial parts of plant specifically stems and leaves were dried in the standard position so that they were protected from light, humidity and infection; also, the samples of suspected to pest and immature organs were separated. All of the samples were powdered using a mill (Model PX-MFC90D) with the mesh size of 1mm for the following analysis. Meanwhile, some of the ecological factors of the sampling area are as follows: the climate of the area is cold, the mean annual rainfall and temperature were 549 mm and 10.1 °C respectively. The average of altitude of area is between 2676–2716 m and the soil texture of area is moderate, semi heavy to heavy.

2.3. Isolation of the essential oils

The air dried of the aerial parts of the plant (100 g) was separated for hydrodistillation using a Clevenger-type apparatus (European pharmacopeia, 1998). The distillation was done for 4 hours. The essential oil obtained was kept in a dark glass sealed vial at 4°C for further analysis. The yields of the oils were calculated based on dried weight of the plant material.

2.4. Gas chromatography/mass spectrometry (GC/MS) analysis

The oils were examined by GC/MS, using Agilent Technologies GC/MS 7890A (built on the industry-leading Agilent GC platform, USA) in the central laboratory of Isfahan University of Technology (IUT), Iran. The injector temperature was set at 220 °C for 5 minutes with a split ratio of 1:10. A 1 µl and a volume of 1000 ppm of the obtained oil solution in a GC grade (*n*-hexane) were injected. To isolate different oil components, a linear temperature program was adjusted as described here: first of all, the column was kept at 50 °C for 2 minutes, ramped at a rate of 10 °C/min to 150 °C, and held isothermal for 5 minutes; then the ramp of 20 °C/min was applied up to 220 °C and held isothermal for 10 minutes. The mass spectrometry detector was taken at 70 eV and the mass range was 40–400 *m/z*. The identification of components was carried out using their recorded spectra by utilizing the spectrometric electronics libraries provided by the instrument software (Wiley, 1994; Adams, 2001). Also, the comparative amounts of the essential oil constituents were stated and recorded as percentage obtained via retention time (*R_i*) and peak area (Area%).

3. Results and discussion

The light yellowish essential oil of *Dorema ammoniacum* D. Don. was obtained with the yield of 0.31 % (v/w) based on the dried weight of the aerial parts. The major compounds of the essence in dried organs were formed about 67% of the total oils so that mentioned identified compounds were Limonene (49.18%), Neophytadiene (10.21%), beta Caryophyllene (3.54%), Phytol (2.61%) and Neryl acetone (1.46%). In this study, 89 compounds belonged to seven main chemical groups consists of nitrogen containing compounds such as terpenes, hydrocarbons, alcohols, ketones and others compounds were identified in the essence of *Dorema ammoniacum* D. Don. Some of these groups were divided to more minor chemical functional

groups for example: nitrogen containing compounds including (amines, alkaloids and amides), terpenoids (monoterpenes, sesquiterpenes and diterpenes), hydrocarbons (aliphatic, aromatic and cyclic hydrocarbons) and the others of remaining groups were assigned in the next levels. The quantitative amounts of mentioned compounds in order of priorities were as follows: nitrogen containing compounds including amines (73.5 %), alkaloids (17.2 %) and amides (9.3 %), terpenoids consists of monoterpenes (82.88 %), sesquiterpenes (12.97 %) and diterpenes (4.14 %), hydrocarbons involving aliphatic hydrocarbons (88.66 %), aromatic hydrocarbons (7.5 %), cyclic hydrocarbons (3.83 %), alcohols (2.76 %), ketones (3.95 %) and miscellaneous compounds (2.22 %). The specifications of the entire identified constituents of the essential oils of *Dorema ammoniacum* D. Don. along with their retention time and also GC/MS chromatogram of it are shown in Table 1 and figure 2 (Table 1 and Fig 2).



Fig 1. *Dorema ammoniacum* D. Don.

In the current research, two major groups of constituents namely nitrogen containing compounds and terpenes were reported more than the other compounds. Overall, nitrogen compounds play an important role in many aspects of life and commercial processes. Nitrogen is a molecule in every major drug class in pharmacology and medicine (Bussau, 1999; Tamras, 1995). On the other hand, terpenes are one of the large group of chemical compounds with more minor divisions which sometimes called isoprenoids. These compounds are used extensively for their aromatic

qualities and play a role in traditional herbal remedies (Connolly and Hill, 1996; Mann et al., 1994). In this regard, a set of concise studies in relation to identification of constituents of *Dorema ammoniacum* D. Don. were carried out around the world. In most of which existence of different Terpenoids compounds have been confirmed and some of the mentioned compounds were similar to the ones identified in this report (Mansouji Avval et al., 2013; Yousefzadi et al., 2011b; Yassa et al., 2014; Batooli et al., 2014). In spite of the reported constituents in the previous studies, there were a wide range of new identified compounds which were not reported in any of the previous research.




Fig 2. GC/MS chromatogram of *Dorema ammoniacum* D. Don.

The findings of the present research shows similarities and differences of essence constituents in mentioned organs of *Dorema ammoniacum* D. Don. compared to the previous conducted research. There are two significant reasons for such variances including, genetic characteristics (different genotypes of plants) and special environmental conditions (Omidbeigi, 2000).

Table 1. The specifications of the entire identified constituents of the essential oils of *Dorema ammoniacum* D. Don.

No.	Compounds ^a	Rt ^b	%
1	2,5-Dimethyltetrahydrofuran	3.162	0.25
2	Methylbenzene	3.222	0.10
3	Cyclohexane, 1,3-dimethyl, <i>trans</i>	3.351	0.3
4	Valeraldehyde	3.456	0.12
5	Octane	3.546	0.65
6	Imidazole-5-carboxylic acid, 2- amino	3.652	0.11
7	Butyl acetate	3.848	0.33
8	Cyclohexane, 1,2-dimethyl, <i>cis</i>	4.149	0.36
9	N,N-dimethyl cyclobutane-1,1- bis(methylamine)	4.194	0.06
10	Furfural	4.300	0.44
11	Deuteroacetone	4.353	0.18
12	Styrene	5.249	0.18
13	1-Methyldodecylamine	5.317	0.11
14	α -Pinene	5.882	0.32
15	Aziridine, 1-(2-buten-2-yl)	6.402	0.47
16	Sabinene	6.541	0.24
17	β -Myrcene	6.824	0.26
18	Decane	6.937	0.52
19	Acetamide, 2-chloro	7.209	0.08
20	Limonene	7.585	49.18
21	2-Butylamine	7.751	0.31
22	2-Pyridinepropanoic acid, α - methyl- β -oxo- ethyl ester	7.819	0.24
23	Phenyl Acetaldehyde	8.007	0.15
24	2-Amino-4-hydroxypteridine-6- carboxylic acid	8.520	0.07
25	Benzenamine, N-methyl-2-(2- propenyl)	8.572	0.17
26	Undecane	8.610	0.5
27	α -Terpinolene	8.723	0.61
28	Aramine	9.175	0.29
29	Limonene oxide	9.409	1.26
30	Isopropyl isocyanate	9.718	0.45
31	2-Hexanamine, 4-methyl	9.914	0.63
32	2,6-Dimethyl-4-pyrone	10.095	0.27
33	Dodecane	10.162	0.97
34	2,4-Hexadiene, 3-methyl	10.275	0.07
35	β -Fenchyl alcohol	10.351	1.27
36	Homarine	10.404	0.48
37	2-(N,N-dimethyl hydrazino) cyclohexane carbonitrile	10.494	0.19
38	Phenol, 4-(2-aminopropyl)	10.554	0.5
39	2-Oxo-3-methyl-cis-perhydro- 1,3-be nzoazine	10.607	0.35
40	<i>trans</i> -Carveol	10.697	0.58
41	Benzene, (2-fluoro-2- methoxycyclopropyl)	10.826	0.35
42	2(1H)-Naphthalenone, octahydro-8a-methyl, <i>trans</i>	10.908	1.14
43	p-Methoxyamphetamine	11.044	0.57
44	2-Cyclohexen-1-one, 2-methyl- 5-(1-methylethenyl)	11.127	0.32
45	2-Methyl amphetamine	11.255	0.13

46	Tridecane	11.617	0.55
47	1-(3,5-Dimethyl-1-adamantanoyl) semi carbazide	11.767	0.52
48	Benzenemethanol, alpha-(1-aminoethyl)	12.084	0.05
49	Naphthalene, 1,2-dihydro-1,1,6-trimethyl	12.672	0.25
50	Rimantadine	12.853	0.1
51	Heptadecane	12.981	0.36
52	β-Caryophyllene	13.501	3.54
53	2-Amino-1-(O-methoxyphenyl) propane	13.591	0.35
54	Calarene	13.651	0.43
55	Neryl acetone	13.734	1.46
56	2-Methoxy amphetamine	14.164	0.63
57	1-[a-(1-Adamantyl)benzylidene] thiosemicarbazide	14.405	0.23
58	Benzenemethanol, 3-hydroxy-alpha-[(methylamino) methyl] delta-Cadinene	14.465	0.2
59	delta-Cadinene	14.699	1.11
60	1 S-cis-Calamenene	14.797	0.16
61	Aptrol	14.857	0.35
62	Benzenemethanol, 4-hydroxy-alpha-[(methylamino) methyl] Nerolidol	14.925	0.31
63	Nerolidol	15.128	0.36
64	N-Methyl-propylamine	15.384	0.11
65	Pentadecane	15.475	0.32
66	Farnesol 2	15.513	1.37
67	Caryophyllene oxide	15.618	1.36
68	2-Hexanamine, 5-methyl	16.145	0.1
69	Benzenemethanol, 2-(2-aminopropoxy)-3-methyl	16.274	0.22
70	2-Propanamine	16.703	0.1
71	2-Aminononadecane	16.922	0.27
72	Methanimidamide, N,N-dimethyl-N-phenyl	17.035	0.34
73	Methylpent-4-enylamine	17.713	0.7
74	Cyclobutanol	18.014	0.3
75	Neophytadiene	18.105	10.21
76	Hexahydro farnesyl acetone	18.218	0.86
77	n-Hexylmethylamine	18.293	0.06
78	Ethylene bromohydrin	18.361	0.46
79	Didodecyl phthalate	18.549	0.21
80	3-Propoxyamphetamine	18.836	0.09
81	Glycine, N-(N-acetyl-L-alanyl) butyl ester	19.069	0.2
82	2-Propenamide	19.250	0.13
83	2-Heptanol, 6-amino-2-methyl	19.348	0.05
84	Benzene ethanamine, 4-fluoro-beta,3-dihydroxy-N-methyl	19.484	0.08
85	1H-Indole-3-ethanamine, 6-fluoro-beta-methyl	19.529	0.05
86	Nonadecane	20.689	0.27
87	Phytol	20.848	2.61
88	Eicosane	22.460	0.83
89	Cyclotrisiloxane, hexamethyl	24.480	0.62
	Total identified		99.01

^a Compounds listed in order of elution on the DB5 column.

^b Retention time (as min)

Nevertheless, other factors such as phenology stages, the period of growing season and different organs of plants have been taken into consideration. It should be noted, our efforts have been focused on investigation and introduction of the new major chemical compounds which would be concisely described by the end of this study. The molecular structure of the major chemical compounds from the essential oil of *Dorema ammoniacum* D. Don. is shown in figure 3 (Fig. 3).

Limonene is the first compound that is widely used in the process of manufacturing and production of various cosmetic products. Numerous applications are mentioned for this compound in literatures. Some of them are as follows: air care products, cleaning and furnishing care products, laundry and dishwashing products, personal care products and it has been considered as biofuel (Connolly and Hill, 1996; Mann et al., 1994; Bauer, et al., 2001; Ming Lin et al., 2015).




Fig 3. The molecular structure of the major chemical compounds from the essential oil of *Dorema ammoniacum* D. Don.

Neophytadiene is the second compound. It is one of the major compounds of *Nicotiana tobacum* L. that exists in leaves, stems and wastes of mentioned plant or is created by the hydration of phytol. It is a potent antimicrobial and anti-inflammatory compound and in some of literatures reviews it is identified as an antifungal terpenoid in the red alga, *Centroceras clavulatum* (C. Agardh) and Montagn. Other applications are as listed: antipyretic, analgesic and

vermifugic, which cures sores and inflammation (Stojanovic *et al.*, 2000; Mendiola *et al.*, 2008; Carretero *et al.*, 2008). The third compound is beta Caryophyllene. This compound is extensively seen in the essential oils of various plants since it has several biomedical properties. The mentioned compounds contributes in the odor and flavor of the plants and also is used in lotions, creams and soaps (Skold *et al.*, 2006). Moreover, several therapeutic features of this compound have been reported in literatures. Some of the most important of them are as follows: anti-anemic effect, anti-inflammatory agent and anti-mutagen, a protector of gastric injury and oxidative stress and finally anti-carcinogen (Legault and Pichette, 2007; Di Sotto *et al.*, 2008; Da Silva *et al.*, 2007). Phytol is the fourth compound that like limonene are used in the fragrance industry and it could be used for manufacturing combined forms of the vitamins E and K₁ (Daines *et al.*, 2003; Mc Ginty *et al.*, 2010). The latest major compound is beta Neryl acetone. This compound has been reported as one of the major combination in some angiosperm (flowering plant) families such as Rutaceae, Burceraceae, Poaceae, Lamiaceae, Lauraceae, Verbenaceae, Asteraceae, Cuppressaceae and Hypericaceae (Pengely, 1996). It is classified in the group of flavor and fragrance agent and participates in the manufacture of various solvents. Additionally, therapeutics effects of the mentioned compound in some studies related to medical field has been proven (Curtis and Williams, 2001; Curzons *et al.*, 1999; Huttenlocher *et al.*, 1971; Gilbert *et al.*, 2000).

4. Conclusion

In this study, we have investigated existing constituents in the extract of *Dorema ammoniacum* D. Don. in one of its habitats in the Central Zagros, Iran. It led to discovery of some new constituents for the first time which were not reported in the previous studies. As discussed in the results section two major factors namely genetic features and environmental conditions have a significant contribution in shaping up and appearance of the various constituents. However, there are some habitats of the mentioned plant on earth that no phytochemical studies have been done on them and it needs to be further researched. *Dorema ammoniacum* D. Don. due to therapeutic properties is used in Iran folk medicine and it has various applications in cosmetics,

detergent industries and food industry. Also, in the contemporary medicine the resin of mentioned plant has been utilized for the treatment of asthma and its case study is still ongoing. Finally, Iran due to its abundant species richness and particular climate conditions can be regarded as a unique country in the world in terms of medicinal species.

5. Acknowledgments

The authors appreciatively acknowledge from Department of Natural Resources and Nanotechnology and Advanced Materials Institute at Isfahan University of Technology for providing facilities and equipment required for the research. Authors are also thankful to the central laboratory of the University for helping to conduct of GC/MS analysis.

6. References

- Adams, R.P. 2007. *Identification of Essential oils Components by Gas Chromatography/Mass Spectroscopy*, fourth ed. Allured Publishing Corporation, Carol Stream, IL, USA.
- Amin, G.R. 1991. *Popular medicinal plants of Iran*. Iranian Research Institute of Medicinal Plants Iran. PP 262.
- Appendino, G., Nano, G.M., Viterbo, D., Munno, G.D., Cisero, M., Palmisano, G. and Aragno, M. 1991. Ammodoremin, a epimeric mixture of prenylated chromandiones from ammoniacum. *Helvetica Chimica Acta*, 74: 495–500.
- Arnone, A., Nasini, G., De Pava, O.V. and Cammarda, L. 1991. Isolation and structure elucidation of doremine A, a new spiro-sesquiterpenoidic chroman-2,4-dione from ammoniac gum resin. *Gazzetta Chimica Italiana Journal*, 121: 383–386.
- Batooli, H., Haghiri Ebrahimabadi, A., Mahmodi, B. and Mazochi, A. 2014. Comparison of essential oil chemical composition of vegetative and reproductive organs of *Dorema ammoniacum* D. Don. in Shahsavaran Kashan, *Iranian Journal of Medicinal and Aromatic Plants*, 30: 746–755.
- Bauer, K., Garbe, D., and Surburg, H. 2001. *Common fragrance and flavor materials*, 4th Ed. Wiley–VCH verlag GmbH, Germany.
- Bown, D. 1995. *Encyclopedia of Herbs and Their Uses*. Dorling Kindersley, UK.

- British Herbal Pharmacopoeia. 1983. The British Herbal and Medical Association, West Yorkshire, UK. PP 24.
- British Pharmacopoeia. 1993. Vol. 2, HMSO, London. UK, PP. A165–169.
- Bussau, A. 1999. *Soluble Nitrogen Containing Compounds in Eucalyptus Globulus (Labill.) Foliage: Effect of Site, Season and Nutrient Addition*. University of Melbourne, Australia. PP 244.
- Carretero, M.E., Lopez Perez, J.L., Abad, M.J., Bermejo, P., Tillet, S., Israel, A. and Noguera, P.B. 2008. Preliminary study of the anti-inflammatory activity of hexane extract and fractions from *Bursera simaruba* (Linneo) Sarg, (Burseraceae) leaves. *Journal of Ethnopharmacology*, 116: 11–15.
- Chevallier, A. 2001. *Encyclopedia of Medicinal Plants*. Dorling Kindersley, UK. PP 285.
- Connolly, J.D., and Hill, R.A. 1996, *Terpenoids*. Chappman and Hall Company, UK, PP. 291–356.
- Convention on the Elaboration of a European Pharmacopoeia. 1998. European Pharmacopoeia: Supplement. Council of Europe.
- Curtis, T., and Williams, D.G. 2001. *Introduction to Perfumery*. Micelle Press, USA.
- Curzons, A.D., Constable, D.C. and Cunningham, V.L. 1999. Solvent selection guide: a guide to the integration of environmental, health and safety criteria into the selection of solvents. *Clean Products and Processes*, 1: 82–90.
- Daines, A., Payne, R., Humphries, M. and Abell, A. 2003. The Synthesis of Naturally Occurring Vitamin K and Vitamin K Analogues. *Current Organic Chemistry*, 7: 1625–1634.
- Da Silva, S.L., Figueiredo, P.M. and Yano, T. 2007. Chemotherapeutic potential of the volatile oils from *Zanthoxylum rhoifolium* Lam leaves. *European Journal of Pharmacology*, 576: 180–188.
- Di Sotto, A., Evandri, M.G. and Mazzanti, G. 2008. Antimutagenic and mutagenic activities of some terpenes in the bacterial reverse mutation assay. *Mutation Research*, 635: 130–133.
- Evans, W.C. 1996. *Trease and Evans' Pharmacognosy*. 14th ed.; W.B. Saunders Company, UK.
- Farhang, H.R., Vahabi, M.R. and Allafchian, A.R. 2015. Chemical compositions of the essential oil of *Gundelia tournefortii* L.(Asteraceae) from Central Zagros, Iran, *Journal of Herbal Drugs*, 6: 227–233.
- Gilbert, D.L., Pyzik, P.L. and Freeman, J.M. 2000. The ketogenic diet: seizure control correlates better with serum beta-hydroxybutyrate than urine ketones. *Journal of Child Neurology*, 15: 787–790.
- Grieve, M. 1971. *A Modern Herbal*. Vol. 1, Peter Smith Publisher, USA.
- Howes, F. 1950. Age-old resins of the Mediterranean region and their uses. *Economic Botany*, 4: 307–316.
- Huttenlocher, P.R., Wilbourn, A.J. and Signore, J.M. 1971. Medium chain triglyceride as a therapy for intractable childhood epilepsy. *Neurology*, 21: 1097–1103.
- Irvani, N., Solouki, M., Omidi, M., Zare, A.Z. and Shahrazi, S. 2010. Callus induction and plant regeneration in *Dorema ammoniacum* D. an endangered medicinal plant. *Plant Cell Tissue and Organ Culture*, 100: 293–300.
- Langenheim, J.H. 2003. *Plant resins: chemistry, evolution, ecology, and ethnobotany*. Timber Press, USA.
- Legault, J. and Pichette, A. 2007. Potentiating effect of β -caryophyllene on anticancer activity of α -humulene, isocaryophyllene and paclitaxel. *Journal of Pharmacy and Pharmacology*, 59: 1643–1647.
- Mann, J., Davidson, R.S., Hobbs, J.B., Banthorpe, D.V. and Harborne, J.b. 1994. *Natural products: Their Chemistry and Biological Significance*. Longman Group, UK.
- Mansouji Avval, M., Sajjadifar, S. and Sadeghei Takallo, M. 2013. Chemical Composition of the Essential Oils from Flowers, Stems and Roots of *Dorema ammoniacum* D.Don. from Iran. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 4: 640–644.
- Mc Ginty, D., Letizia, C.S. and Api, A.M. 2010. Fragrance material review on phytol. *Food and Chemical Toxicology*, 48: 59–63.
- Mc Lafferty, F.W. 2013. *Wiley registry of mass spectral data*. tenth ed. Mass Spectrometry Library Search System Bench-Top/PBM, New York, USA.
- Mendiola, J.A., Santoyo, S., Cifuentes, A., Reglero, G., Ibanez, E. and Senorans, F.J. 2008. Antimicrobial activity of sub- and supercritical CO₂ extracts of the green alga *Dunaliella salina*. *Journal of Food Protection*, 71: 2138–2143.
- Ming Lin, F., Neil, E., Marsh, G. and Nina Lin, X. 2015. Recent progress in hydrocarbon biofuel synthesis:

- pathways and enzymes. *Chinese chemical letters*, 26: 431–434.
- Mozaffarian, V.A. 1996. *Dictionary of Iranian Plant Names*. Farhang Moaser Publication Iran, PP. 145.
- Omid Beigi, R. 2000. *The production and processing of medicinal plants*. Fekr Ruose Publication, Iran.
- Parsa, A. 1959. Medicinal plants and drugs of plant origin in Iran III. *Plant Food for Human Nutrition*, 6: 137–156.
- Pengely, A. 1996. *The constituents of medicinal plants*. Allen and Unwin, Australia.
- Rajani, M., Saxena, M.N., Ravishankara, N., Desai, N. and Padh, H. 2002. Evaluation of the Antimicrobial Activity of Ammoniacum Gum from *Dorema ammoniacum*. *Pharmaceutical Biology*, 40: 534–541.
- Rechinger, K. H. 1980. *Flora Iranica*. Umbelliferae; Akademische Druck- u. Verlagsanstalt Graz, Austria. PP. 234.
- Sajjadi, S.E., Ghassemi, N. and Mohammadzamani, P. 2007. Chemical Constituents of the Essential Oil of *Dorema ammoniacum* D. Don. Leaf, an Iranian Resinous Plant. *Revue des regions Arides*, 1: 194–197.
- Skold, M., Karlberg, A.T., Matura, M. and Borge, A. 2006. The fragrance chemical β -caryophyllene air oxidation and skin sensitization. *Food and Chemical Toxicology*, 44: 538–545.
- Stojanovic, G., Palic, R., Alagic, S. and Zekovic, Z. 2000. Chemical composition and antimicrobial activity of the essential oil and CO₂ extracts of semi-oriental tobacco *Otlja*. *Flavour and Fragrance Journal*, 15: 335–338.
- Tarmas, S. 1995. *Electrophilic Aromatic Thallation of Nitrogen containing Compounds*. California State University, Fresno, USA.
- Worwood, V.A. 1991. *The Complete Book of Essential Oils and Aromatherapy*. New World Library Publication California, USA.
- Yassa, N., Rustaie, A., Tavakoli, S., Delnavazi, M.R. and Batooli, H. 2014. Antioxidant and antibacterial activities of the essential oils and extracts of *Dorema ammoniacum* roots and aerial parts, *Research Journal of Pharmacognosy*, 1: 11–18.
- Yousefzadi, M., Heidari, M., Akbarpour, M., Mirjalili, M.H., Zeinali, A. and Parsa, M. 2011a. In vitro Cytotoxic Activity of the Essential Oil of *Dorema ammoniacum* D. Don. *Middle East Journal of Scientific Research*, 7: 511–514.
- Yousefzadi, M., Mirjalili, M.H., Alnajjar, N., Zeinali, A. and Parsa, M. 2011b. Composition and *in vitro* antimicrobial activity of the essential oil of *Dorema ammoniacum* D. Don. fruit from Iran. *Journal of Serbian Chemical Society*, 76: 857–863.
- Zarshenas, M.M., Arabzadeh, A., Tafti, M.A., Kordafshari, G., Zargarani, A. and Mohagheghzadeh, A. 2013. Application of Herbal Exudates in Traditional Persian Medicine. *Galen Medical Journal*, 1: 78–83.