



The Comparison of intravenous and oral administration of anise plant on BALB/c mice with listeriosis

Amin Hashempour^{1*}, Mohammad Hosein Sadeghi Zali², Saber Yousefi³

¹Young Researcher and Elites club, Urmia Branch, Islamic Azad University, Urmia, Iran;

²Department of Bacteriology, Veterinary Faculty, Islamic Azad University, Urmia Branch, Urmia, Iran;

³Department of Microbiology, Virology and Immunology, Urmia University of Medical Sciences, Urmia, Iran;

ARTICLE INFO

Type: Original Research

Topic: Medicinal Plants

Received February 19th 2016

Accepted April 15th 2016

Key words:

- ✓ *Pimpinella anisum*
- ✓ Extract
- ✓ Essential oil
- ✓ *Listeria monocytogenes*
- ✓ Antibacterial

ABSTRACT

Background & Aim: Anise studies have demonstrated different properties such as anti-cancer, anti-fungal, antibacterial, antioxidant, strengthen the liver, gastrointestinal tract, wounds heal and strengthen the nerve cells. Outbreaks of listeriosis in the general population are 7.0 per 100,000 populations. Pregnant women, infants, the elderly and immunocompromised patients or those weaknesses comprise most cases. The aim of the present study was to investigate the effect of intravenous and oral administration of anise herb to BALB/c race of mice with listeriosis.

Experimental: 2 units of *listeria monocytogenes* were prepared in 0.5 Mc Farland concentrations and injected to BALB/c trace mouse intraperitoneal. Before and after treatment, the mice were examined for agglutination test.

Results: The results of this study indicated that the effect of oral treatment with aqueous extract is better than injection of essential oil of anise.

Recommended applications/industries: According to results of the present study, anise aqueous extract was recommended as a potential antibacterial agent against *listeria monocytogenes*.

1. Introduction

In recent years much attention has been devoted to natural antioxidant and their association with health benefits (Arnous et al., 2001; Huda-Faujan et al., 2009). Plants are a large source of new bioactive molecules with therapeutic potentials. Only a small percentage of living plants on Earth have been phytochemically investigated. Plants are thus an enormous reservoir of pharmaceutically valuable molecules to be discovered (Hostettmann et al., 1998; Balandrin et al., 1985; Kosalec et al., 2005).

Pimpinella anisum L. (Apiaceae) is an annual herb indigenous to the near east and widely cultivated in the Mediterranean regions as well as Mexico and Chile. It has been used as an aromatic herb and spice since Egyptian times (Kosalec et al., 2005). The star anise and anise have both been widely used in Iranian traditional medicine for their antimicrobial effects (Amin, 2005). The antimicrobial properties of these species such as anise and star anise have been reviewed by several researchers (Kosalec et al., 2005; Gulcin et al., 2003; Sungog et al., 2007; Yazdani et al. 2008).

Anise seeds are used as analgesic in migraine and also as carminative, aromatic, disinfectant, and diuretic in traditional medicine (Amin et al., 2005). Aniseed has

warm and dry nature and can increase milk production, menstruation, urine, and sweat secretion and also making good complexion. It is also effective in polishing of teeth. In some traditional texts, anise is mentioned for melancholy, nightmare, and also in treatment of epilepsy and seizure (Mirheydar *et al.*, 2001; Aghili Khorasani *et al.*, 2001).

Aniseed contains 1.5–6.0% of a volatile oil consisting primarily of trans-anethole and also as much as 8–11% of lipids rich in fatty acids, such as palmitic and oleic acids, as well as approximately 4 % of carbohydrates, and 18 % of protein (Besharati-Seidani *et al.*, 2005). Other studies have demonstrated the presence of eugenol trans-anethole, methylchavicol, anisaldehyde, estragole, coumarins, scopoletin, umbelliferone, estrols, terpene hydrocarbons, polyenes, and polyacetylenes as the major compounds of the essential oil of anise seed (Gulcin *et al.*, 2003). Study of the essential oil of *Pimpinella anisum L.* fruits by GC and GC-MS showed the presence of trans-anethole (93.9%) and estragole (2.4%).

Other compounds that were found with concentration higher than 0.06% were (E)-methyleugenol, α -cuparene, α -himachalene, β -bisabolene, p-anisaldehyde, and cis-anethole (Ozcan and Chalchat, 2006). In another study for determination of the composition of essential oil of *Pimpinella anisum L.* fruits obtained from different geographical areas of Europe, in addition to the major components (trans-anethole (76.9–93.7%) and γ -himachalene (0.4–8.2%), some other compounds such as trans-pseudoisoeugenyl 2-methylbutyrate, p-anisaldehyde, and methylchavicol were also identified in essential oil (Orav *et al.*, 2008).

Essential oil and methanol extract of these plants exhibited antibacterial activity against most tested pathogens, and the maximum effect was observed against *Staphylococcus aureus*, *Bacillus cereus*, and *Proteus vulgaris*. However, combination of essential oil and methanol extracts of these plants showed an additive effect against most tested bacteria especially *Pseudomonas aeruginosa* (Al-Bayati, 2008). The antibacterial potential of aqueous decoctions of black pepper, bay leaf, aniseed, and coriander against 176 bacterial isolates belonging to 12 different genera were detected by the mean of disc diffusion technique. The findings showed that the aqueous decoction of black pepper was the most bacterial-toxic exhibited 75% antibacterial activity and decoction of aniseed exhibited

18.1% antibacterial activity (maximum antibacterial activities exhibited against *Micrococcus roseus*) (Chaudhry & Tariq, 2006). In addition to antibacterial activity, the essential oil of aniseed showed significant inhibitory activity against fungi, and the most active component of it was anethol (Shukla & Tripathi, 1987).

2. Materials and Methods

2.1. Preparation of mice

128 mice were prepared from research center of Zist Faravarde Pars in Rasht city, Guilan. Mice were transferred to animal houses of Islamic Azad University, Urmia branch.

2.2. Bacteria injection

After sub culture, bacteria transferred to peptone water media. After 24h incubation L.P was injected to mice.

2.3. Agglutination test after bacteria injection

Listeria Agglutination kits were prepared from research center of Zist Faravarde Pars in Rasht city, Guilan. Agglutination test was done 3 days after injection to ensure of listeriosis in mice. Blood samples were taken from tail vein.

Table 1. Characteristics of bacteria

S. No	Name	Type	PTCC No
1	<i>Listeria monocytogenes</i>	Gram positive bacteria	PTCC1298

2.4. Treatment by injection

After ensuring prevalence of listeriosis in mice subcutaneous injection were done daily at a rate of 1 unit by insulin syringes.

2.5. Treatment with anise seed extract in oral way

The extract was fed to mice by 5g per 100ml instead of drinking water.

2.6. Agglutination test after treatment

After one week of injection of anise essential oil, agglutination test was done to ensure that the treatment is completed.

3. Results and discussion

Anise (*Pimpinella anisum* L), is an annual important spice and medicinal plant belonging to the family of *Apiaceae*, and native to Mediterranean region. Today, anise seeds are important natural raw materials which are used in pharmaceuticals, perfumery, food and cosmetic industries (Ross, 2001). In this study the effect of aqueous extracts and essential oils were compared and it was found that the effect of aqueous extract applied orally was more than injection of essential oil.

3.1. Agglutination test before treatment

Agglutination test was performed on 128 rats and almost the incidence of listeriosis was 100% in mice. Only four rats were not infected and therefore kept separately.

3.2. Treatment

According to table 2, treatment by injection with essential oils and orally water extracts was done on 50 mice separately for a week. According to results, antibacterial effect by injection of essential oils and orally water extracts were 24% and 74%, respectively.

Table 2. Number of cured and not-cured mice treated with anise plant essential oil and extract in 7 days.

Total samples	Treatment	Cured	not-cured
50	Water extract (oral)	37	13
50	Essential oil (Subcutaneous)	12	38
24	Control	0	24
	Total	49	75
		124	

In many studies antibacterial effect of anise seed was reported. For example, Al-Bayati (2008) was shown antibacterial activity of *Pimpinella anisum*. In another research conducted by Yazdi et al. (2014) antibacterial effect and strengthening immune responses of *Pimpinella anisum* was confirmed in broiler chicks (Yazdi et al., 2014).

Antimicrobial effects of water and ethanolic extracts of anise seeds against 10 bacterial species and *Candida albicans* were studied by Gulcin et al. (2003) with disc diffusion method. They reported that ethanolic extract showed significant inhibitory activity against all tested bacteria but was not effective on *Candida albicans*.

However, the antimicrobial effect of water extract was not detected against gram-negative bacteria, *Pseudomonas aeruginosa*, and *Escherichia coli*, but it was effective against *Candida albicans* (Gulcin et al., 2003).

Also, alcoholic extracts of *Pimpinella anisum* seeds showed antibacterial activity against *Micrococcus luteus* and *Mycobacterium smegmatis* (Ates & Erdogru, 2003).

The antibacterial activities of the aqueous, methanol, acetone and petroleum ether extracts of *Pimpinella anisum* L. fruits were tested against 4 pathogenic bacteria (*Staphylococcus aureus*, *Streptococcus pyogenes*, *Escherichia coli*, and *Klebsiella pneumoniae*) by disc diffusion method. The results showed that only aqueous and methanol extracts exhibited fair antibacterial activity against all of the tested bacteria and the aqueous extract was found to be more effective than methanolic extract, whereas acetone and petroleum ether extracts did not show inhibitory effect on growth of the pathogenic tested bacteria (Akhtar et al., 2008).

4. Acknowledgment

Authors are thankful to Young Researcher and Elite club (Islamic Azad University Urmia branch) for providing financial assistance in the form of a major research project. Special thanks to Dr. Ashkan Khoda bandeloo, Dr. Haleh Kangarlou, Dr. Mahmoud pour Yousef, Dr. Pezhman Mohamadi, Mr. Reza Delshad, Mr. Amin khorsandian and Mr. Mostafa Ghorban zadeh for helping us in carrying out experiments.

5. References

- Aghili, M.H. 2001. Khorasani, Makhzan al Advieh, Bavardaran Press. Research institute for Islamic and Complementary Medicine, Iran University of Medical Sciences, Tehran, Iran.
- Akhtar, A. A. Deshmukh, A. V. Bhonsle .2008. "In vitro Antibacterial activity of *Pimpinella anisum* fruit extracts against some pathogenic bacteria," *Veterinary World*. 1(9): 272–274.
- Al-Bayati, F. A. 2008. Synergistic antibacterial activity between *Thymus vulgaris* and *Pimpinella anisum* essential oils and methanol extracts. *Journal of Ethnopharmacology*. 116(3): 403–406.

- Amin, G. 2005. Popular medicinal plants of Iran. Vice-Chancellorship of Research, Tehran University of Medical Science Press, Tehran, Iran.
- Arnous A, Makris DP, Kefalas P. 2001. Effect of principal polyphenolic components in relation to antioxidant characteristics of aged red wines. *Journal of Agriculture and Food Chemistry*. 49: 5736-5742.
- Ates D. A and O. T. Erdogru. 2003. Antimicrobial activities of various medicinal and commercial plant extracts. *Turkish Journal of Biology*, 27: 157–162.
- Balandrin, M. F., Klocke, J. A., Wurtele, E. S. and Bollinger, W. H. 1985. Natural plant chemicals: sources of industrial and medicinal materials, *Science*. 228: 1154–1160.
- Besharati-Seidani, A. Jabbari, and Y. Yamini. 2005. Headspace solvent microextraction: a very rapid method for identification of volatile components of Iranian *Pimpinella anisum* seed. *Analytica Chimica Acta*. 530(1): 155–161.
- Chaudhry N. M. and Tariq. P. 2006. Bactericidal activity of black pepper, bay leaf, aniseed and coriander against oral isolates. *Pakistan Journal of Pharmaceutical Sciences*. 19(3): 214–218.
- Gulcin I, Oktay M, Kirecci E, Ku freviog Lu OI. 2003. Screening of antioxidant and antimicrobial activities of anise (*Pimpinella anisum* L.) seed extracts. *Food Chemistry*. 83: 371 - 382.
- Hostettmann, K., Potterat, O. and Wolfender J.L. 1998. The potential of higher plants as a source of new drugs. *Chimia*. 52: 10–17.
- Huda-Faujan, N., Noriham, A., Norrakiah, A. S. and Babji, A. S., .2009. Antioxidant activity of plants methanolic extracts containing phenolic compounds, *African Journal of Biotechnology*. 8(3): 484-489
- Kosalec I, Pepeljnjak S, & Kustrak D. 2005. Antifungal activity of fluid extract and essential oil from anise fruits (*Pimpinella anisum* L., apiaceae). *Acta Pharmaceutica*. 55 (4): 377 - 85.
- Mirheydar, H. 2001. Herbal information: usage of plants in prevention and treatment of diseases, Islamic Culture Press Center, Tehran, Iran.
- Orav, A. Raal, and E. Arak. 2008. Essential oil composition of *Pimpinella anisum* L. fruits from various European countries. *Natural Product Research*. 22, 3: 227–232.
- Ozcan, M. M and Chalchat, J. C. 2006. Chemical composition and antifungal effect of anise (*Pimpinella anisum* L.) fruit oil at ripening stage. *Annals of Microbiology*, 56, 4: 353– 358.
- Ross, I.A., 2001. Medicinal plants of the world: chemical constituents, traditional and modern medicinal uses, Volume 2. Humana press, Totowa, New Jersey, pp.363-374.
- Shukla, H. S and Tripathi, S. C.1987. Antifungal substance in the essential oil of anise (*Pimpinella anisum* L.),” *Ultural and Biological Chemistry*, 51, 7: 1991–1993.
- Sungog L, Park IK, Choi GJ, Lim HK, Jang KS, Cho KY. 2007. Fumigant activity of essential oils and components of *Illicium verum* and *Schizonepeta tenuifolia* against *Botrytis cinerea* and *Colletotrichum gloeosporioides*. *Journal of Microbiology and Biotechnology*. 17(9): 1568 - 1572.
- Yazdani D, Rezazadeh Sh, Amin Gh, Zainal Abidin MA, Shahnazi S, Jamalifar H .2008. Antifungal activity of dried extracts of anise (*Pimpinella anisum* L.) and star anise (*Illicium verum* Hook. f.) against dermatophyte and saprophyte Fungi, 8(5): 24-29.
- Yazdi, F.F, Ghalamkari, G. Toghiani, M., Modaresi, M., Landy, N .2014. Anise seed (*Pimpinella anisum* L.) as an alternative to antibiotic growth promoters on performance, carcass traits and immune responses in broiler chicks, *Asian Pacific Journal of Tropical Disease*. 4(6): 447-451.