



## The effects of placement position and corm size of saffron (*Crocus sativus* L.) on stigma and corm yields in Ankara conditions

Neşet Arslan<sup>1</sup>, Arif Ipek<sup>2</sup>, Amir Rahimi<sup>1\*</sup>, Gökhan Ipek<sup>1</sup>

<sup>1</sup>Ankara University, Agriculture Faculty, Field Crops Department, Ankara, Turkey;

\*Email: [emir\\_10357@yahoo.com](mailto:emir_10357@yahoo.com)

<sup>2</sup>Çankırı University, Science Faculty, Biology Department, Çankırı, Turkey;

### ARTICLE INFO

**Type:** Original Research

**Topic:** Agronomy

Received March 1<sup>st</sup> 2012

Accepted June 15<sup>th</sup> 2012

### Key words:

- ✓ Cultivation
- ✓ *Crocus sativus* L.
- ✓ Vertical placement
- ✓ Stigma

### ABSTRACT

**Background & Aim:** Saffron (*Crocus sativus* L.) formerly was important in Turkey. Saffron cultivation has been decreased and it is now only cultivated in three villages in this country. It is triploid and exclusively propagated in a vegetative way by corms. In Turkey, saffron is traditionally planted by placing corms in rows randomly without grading or sorting; which results in placement of corms in vertical, horizontal and reverse in rows at random. The study aimed to find the effect of corm size and position of placement in rows on stigma and corms yield.

**Experimental:** This study was done at the experimental fields of the Department of Field Crops, Faculty of Agriculture, Ankara University, during 2010-2011 using four different corms size and three placement positions in three replications.

**Results & Discussion:** The results revealed that position of corms and their size had significant effect on stigma yield. Whereas, corm yield was influenced by corm size only.

**Recommended applications/industries:** According this study, vertical placement produced more stigma yield than other ones in planting. Vertical placement corms in rows should be practiced to obtain more stigma yield.

### 1. Introduction

Saffron (*Crocus sativus* L.) is known as one of the earliest cultivated plants (Ipek et al., 2009). This plant is an important crop cultivated as the source of its spice for at least 3,500 years. Saffron is a perennial spice and has been spread out in Mediterranean and west of Asia from 10 west to 80 east degrees of geographical longitude, as well as from 30 to 50 north degrees of geographical latitude and up to 1000 meters from sea level (Kafiet al., 2002). Currently, it is being cultivated more or less intensively in Iran, India, Greece, Spain,

Italy, Turkey, France, Switzerland, Israel, Pakistan, Azerbaijan, China, Egypt, United Arab Emirates, Japan, Afghanistan, Iraq and recently Australia (Tasmania) (Nehviet al., 2007). In the Mediterranean region, saffron is also cultivated on a much smaller scale in Italy, Greece and Turkey. In Turkey, it is currently being grown in the villages of Davutobasi in the Safran bolu district of Karabuk province (Vurdu, 1993). Until the first quarter of 20<sup>th</sup> century, Safran bolu region was a growing and trading center for saffron, and the region is named after saffron itself (Gümüşsuyu, 2003).

Saffron is classified into Magnoliophyta division, class Liliopsida and order Asparagales. It is a member of Iridaceae family and *Crocus* L. genus. Iridaceae family includes about 60 genera and 1,500 species. The plants in this family are herbs with rhizomes, corms or bulbs. The *Crocus* genus includes approximately 80 species worldwide (Fernandez, 2004) and 32 of them exist in Flora of Turkey, 18 of them are endemic to Turkey (Arslanet al., 2007). The major components of saffron are crocins, picrocrocin and safranal. Crocins is responsible for the color of saffron, whereas picrocrocin and safranal are responsible for its bitter taste and aroma (De Juan et al., 2009).

This plant is applicable in food, cosmetic and dying industries as well as artistic activities specifically for pharmaceutical uses. This plant has some advantages including less water requirement, optimal growing season according to the farmer's views and great opportunities for job and exchange remunerative. Recently, it has been taken more into consideration due to its beneficial biological effects specifically its anti-cancerous effects (Abdullayev, 2005).

This geophyte plant is a sterile autumn-flowering species (Moraga et al., 2009) and propagates by vegetative reproduction through the formation of daughter corms from the mother corm (Basker & Negbi, 1983). An important factor in saffron-stigma production is the planting of large corms (Arslanet al., 2007). Corms placement positions (vertical, horizontal and reverse) are other probable factor in saffron-stigma production. In recent years, some studies have been made in order to progress saffron cultivation in Turkey. In these studies, the biggest handicap is the lack of corm material.

This study aimed to evaluate the effects of corms size and placement positions on saffron's stigma and reproductive daughter corm yield (propagation) in the field conditions.

## 2. Materials and Methods

The trial was done at the experimental fields of the Department of Field Crops, Faculty of Agriculture, Ankara University, during 2010-2011. The long term outdoors climatic data of the experimental city (Table1) and soil analysis results of the experimental soil samples in the field (Table2) are shown. Saffron corms

obtained from Agricultural Directory of Karabük Province were used as the study material. Corms were classified according to corm perimeter as 6 cm, 7 cm, 8 cm and 9 cm (Table3).

**Table 1.** The long term outdoors climatic data of the experimental city (1926-2008)

| Months    | Rainfal (mm) | Temperature (C°) | Humidity (%) |
|-----------|--------------|------------------|--------------|
| September | 6,8          | 20,4             | 46,4         |
| October   | 29,0         | 14,9             | 59,1         |
| November  | 49,6         | 5,7              | 72,1         |
| December  | 33,2         | 0,9              | 78,0         |
| January   | 33,1         | 0,7              | 76,5         |
| February  | 38,1         | 0,7              | 73,1         |
| March     | 24,5         | 6,4              | 63,0         |
| April     | 39,8         | 12,6             | 57,8         |
| May       | 47,9         | 16,1             | 56,6         |
| June      | 20,5         | 20,1             | 50,5         |
| July      | 8,8          | 23,5             | 45,9         |
| August    | 6,3          | 23,4             | 46,5         |

**Table2.** Soil analyses results of the experimental soil samples in the field before corm sowing

| Depth             | 0 – 20 (cm) | 20 – 40 (cm) |
|-------------------|-------------|--------------|
| PH9               | 7,33        | 7,26         |
| CaCO3 %           | 9,00        | 10,0         |
| Organic Material% | 1,14        | 1,02         |
| Salt %            | 0,063       | 0,076        |
| Sand %            | 39,99       | 35,31        |
| Clay %            | 22,18       | 24,35        |
| Loam %            | 37,83       | 40,24        |
| P2O5 kg/da        | 5,3         | 4,5          |
| K2O kg/da         | 140         | 122          |

**Table 3.** Used corms (kg/d) in different corm perimeter (cm)

| Corm perimeter (cm) | 6 cm          | 7 cm          | 8 cm          | 9 cm          |
|---------------------|---------------|---------------|---------------|---------------|
| Used corm (kg/d)    | 287,40 (kg/d) | 408,52 (kg/d) | 558,61 (kg/d) | 756,67 (kg/d) |

On 2<sup>nd</sup> June 2011, corms were harvested in the plots. The results were subjected to variance analysis and differences among average values were determined statistically.

Corn planting was done on 30<sup>th</sup> September 2010 using randomized complete block and split-plot design with three replications. Planting was performed as 20 x

10 cm of plant spacing with corm sizes as main plots and placement positions (vertical, horizontal and reverse) as subplots. Each plot consisted of three rows with 60 corms due to lack of corm material. Each plot was  $0.6 \text{ m} \times 2 \text{ m} = 1.2 \text{ m}^2$ . No fertilization and irrigation were applied to the experiment. The flowering harvesting was done at November 11<sup>th</sup>2010.

### 3. Results and Discussion

#### 3.1. Corm yield

Corm yield (kg/d) obtained from different corm size (cm), placement positions and interaction between two factors are shown in Table4, Table5 and Table6 respectively (10 d = 1 ha). Corm yield was affected statistically ( $p < 0.01$ ) by corm size. Corm yield with respect to corm perimeter changed between 545 kg/d (6 cm) and 1377 kg/d (9 cm). There was no significant difference between three placement positions. With respect to this factor, the highest corm yield (931 kg/d) was recorded from the horizontal placement while the lowest yield (859.9 kg/d) was obtained from the reverse placement. Interaction between two factors was not significant. With respect to interaction, the highest corm yield (1441 kg/d) was recorded from the combination of vertical placement and 9 cm of corm size while the lowest yield (535 kg/d) was obtained from the combination of horizontal placement and 6 cm of perimeter.

#### 3.2. Stigma Yield

Stigma yield (g/d) obtained from different corm size (cm), placement positions and interaction between two factors are shown in Table4, Table5 and Table6 respectively. Stigma yield was affected statistically ( $p < 0.01$ ) by corm size and placement position ( $p < 0.01$ ). Stigma yield with respect to corm perimeter changed between 91.4 g/da (7 cm) and 530.2 g/d (9 cm). With respect to placement position, the highest yield (329.3 g/d) was recorded from the vertical placement while the lowest yield (213.9 g/d) was obtained from the reverse placement. Interaction between two factors was not significant. With respect to interaction, the highest stigma yield (657.22 g/d) was recorded from the combination of vertical placement and 9 cm of corm size while the lowest yield (62.22 g/d) was obtained from the combination of reverse placement and 7 cm of perimeter.

#### 3.3. Variation of Corm Yield in Size after harvesting

The variation of corm yield (kg/d) in size after harvesting in different corm perimeter (cm) and placement position are presented in Table 7. Corm number per  $1.2 \text{ m}^2$  and corm weight (kg/d) increased sharply with increasing of corm perimeter. Larger corms produced more corm numbers and corm weight (kg/d) than smaller corms. In every corm size, the corm number in reverse placement position was more than other placement position but the corm weight is random.

**Table 4.** The corm yield (kg/d) and stigma yield (g/d) in different corm perimeter (cm).

| Corm perimeter (cm) | Corm yield (kg/d) | Stigma yield (g/d) |
|---------------------|-------------------|--------------------|
| 6 cm                | 545.00 d          | -                  |
| 7 cm                | 738.90 c          | 91.40 b            |
| 8 cm                | 957.40 b          | 231.40 b           |
| 9 cm                | 1377.00 a         | 530.20 a           |

**Table 5.** The corm yield (kg/d) and stigma yield (g/d) in different placement position

| placement positions | Corm yield (kg/d) | Stigma yield (g/d) |
|---------------------|-------------------|--------------------|
| Vertical            | 923.00 -          | 329.30 a           |
| Horizontal          | 931.00 -          | 309.80 ab          |
| Reverse             | 859.90 -          | 213.90 b           |

The results revealed that corm size had significant effect on corm yield. Many literature reports indicate that there is a positive correlation between corm size and regenerated corm number in saffron (McGimpsey et al., 1997; Arslan, 1997; Gümüşsuyu, 2002; Vurduet al., 2002; Deo, 2003; Vurdu and Güneş, 2004; Çavuşoğlu and Erkel, 2005). New regenerated corm number increased when corm perimeter increased. But this increasing rate is low in small corms. McGimpsey et al. (1997) obtained 10.540 kg/d of corm yield five years after planting using 29 g corms as the seeding material. The effect of placement position on corm yield was not significant but reverses planting produced lowest corm yield. Vertical planting can produce more corm yield. Also the corm yield is affected by planting density, some agricultural practices and so on.

**Table 6.** The corm yield (kg/d) and stigma yield (g/d) in different corm perimeter (cm) and placement position (interaction)

| Corm position<br>Corm size | Corm yield (kg/d) |            |         |         | Stigma yield (g/d) |            |         |         |
|----------------------------|-------------------|------------|---------|---------|--------------------|------------|---------|---------|
|                            | Vertical          | Horizontal | Reverse | Average | Vertical           | Horizontal | Reverse | Average |
| 6 cm                       | 547.78            | 535.00     | 552.22  | 545.00  | -                  | -          | -       | -       |
| 7 cm                       | 741.67            | 772.78     | 702.22  | 738.89  | 87,50              | 124,45     | 62,22   | 91.39   |
| 8 cm                       | 962.22            | 988.89     | 921.11  | 957.41  | 243,06             | 280,00     | 171,11  | 231.39  |
| 9 cm                       | 1441.67           | 1425.56    | 1263.89 | 1377.04 | 657,22             | 525,00     | 408,33  | 530.18  |
| Average                    | 923.33            | 930.56     | 859.86  | -       | 329.26             | 309.82     | 213.89  | -       |

**Table 7.** The variation of corm yield (kg/d) in size after harvesting in different corm perimeter (cm) and placement position

| Corm perimeter (cm) | Placement position | Corm yield ( 6 cm < ) |                            | Corm yield ( 6 cm ) |                            | Corm yield ( < 6 cm ) |                            |
|---------------------|--------------------|-----------------------|----------------------------|---------------------|----------------------------|-----------------------|----------------------------|
|                     |                    | Weight (kg/d)         | Number per1 m <sup>2</sup> | Weight (kg/d)       | Number per1 m <sup>2</sup> | Weight (kg/d)         | Number per1 m <sup>2</sup> |
| 6 cm                | Vertical           | 338.3(65%)            | 44.7(36%)                  | 83.9(16%)           | 18.8(15%)                  | 97.8(19%)             | 60.0(49%)                  |
|                     | Horizontal         | 343.3(66%)            | 44.2(39%)                  | 83.9(16%)           | 19.2(17%)                  | 90.0(18%)             | 50.2(44%)                  |
|                     | Reverse            | 355.5(67%)            | 43.1(34%)                  | 70.5(13%)           | 15.2(12%)                  | 104.4(20%)            | 69.4(54%)                  |
| Corm perimeter (cm) | Placement position | Corm yield ( 7 cm < ) |                            | Corm yield ( 7 cm ) |                            | Corm yield ( < 7 cm ) |                            |
|                     |                    | Weight (kg/d)         | Number per1 m <sup>2</sup> | Weight (kg/d)       | Number per1 m <sup>2</sup> | Weight (kg/d)         | Number per1 m <sup>2</sup> |
| 7 cm                | Vertical           | 369.9(51%)            | 39.2(23%)                  | 132.8(18%)          | 22.7(13%)                  | 219.4(31%)            | 110.8(64%)                 |
|                     | Horizontal         | 397.2(53%)            | 40.2(25%)                  | 148.9(20%)          | 25.0(15%)                  | 206.3(27%)            | 97.2 (60%)                 |
|                     | Reverse            | 305.5(46%)            | 35.2(19%)                  | 147.8(22%)          | 25.2(14%)                  | 213.9(32%)            | 125.6(67%)                 |
| Corm perimeter (cm) | Placement position | Corm yield ( 8 cm < ) |                            | Corm yield ( 8 cm ) |                            | Corm yield ( < 8 cm ) |                            |
|                     |                    | Weight (kg/d)         | Number per1 m <sup>2</sup> | Weight (kg/d)       | Number per1 m <sup>2</sup> | Weight (kg/d)         | Number per1 m <sup>2</sup> |
| 8 cm                | Vertical           | 353.3(37%)            | 30.0(14%)                  | 188.9(20%)          | 24.2(11%)                  | 407.2(43%)            | 160.0(75%)                 |
|                     | Horizontal         | 355.0(36%)            | 29.2(14%)                  | 216.4(23%)          | 26.7(12%)                  | 400.0(41%)            | 159.2(74%)                 |
|                     | Reverse            | 233.9(26%)            | 18.6(8%)                   | 218.3(25%)          | 26.4(11%)                  | 439.4(49%)            | 191.3(81%)                 |
| Corm perimeter (cm) | Placement position | Corm yield ( 9 cm < ) |                            | Corm yield ( 9 cm ) |                            | Corm yield ( < 9 cm ) |                            |
|                     |                    | Weight (kg/d)         | Number per1 m <sup>2</sup> | Weight (kg/d)       | Number per1 m <sup>2</sup> | Weight (kg/d)         | Number per1 m <sup>2</sup> |
| 9 cm                | Vertical           | 308.9(24%)            | 21.1(7%)                   | 255.0(20%)          | 24.2(9%)                   | 725.5(56%)            | 232.2(84%)                 |
|                     | Horizontal         | 315.0(24%)            | 20.6(8%)                   | 380.5(29%)          | 33.9(14%)                  | 618.9(47%)            | 191.9(78%)                 |
|                     | Reverse            | 352.8(26%)            | 21.4(7%)                   | 279.4(20%)          | 24.4(9%)                   | 745.5(54%)            | 243.1(84%)                 |

In this study, as no fertilization and irrigation was applied, soil hardening resulted in reduced flower harvest as well as stigma yield. The results indicated that corm size and placement position had significant

effect on stigma yield. Larger corms and vertical planting produced more stigma yield than smaller corms and reverse planting. Arslan et al. (2007) revealed that best result was obtained from 9.00-11.00

cm perimeter of corm than smaller corms. Other researchers indicated that an important factor in stigma yield is the planting of large corms (İpek et al., 2009; Gresta et al., 2008; Vurduet al., 2002). Hosseini et al. (2004) found different results about stigma yield in different studies.

#### 4. Conclusion

The results of our study indicate that both stigma yield and corm yield were affected by corm size while only stigma yield was affected by placement position. In Turkey, saffron is traditionally planted by placing corms in rows randomly without classification which resulted in placement of corms in vertical, horizontal and reverse in rows at random. This study revealed that corm classification should be necessary before planting. Larger corms should be used in order to get more stigma yield and regenerated corms. Small corms should be planted in a different place in order to be enlarged. As results show, vertical placement produced more stigma yield than other ones in planting. Vertical placement corms in rows should be practiced to obtain more stigma yield.

#### Acknowledgements

The author is highly thankful to Prof. Dr. Khalid Mahmood Khalid for helping in preparation of manuscript and analyzing manuscript data statistically.

#### References

- Abdullayev, F. 2005. Medicinal use of saffron (*Crocus sativus* L.). *Clinical Biochemistry*, 38: 865-6.
- Arslan, N., Gürbüz, B., İpek, A., Özcan, S., Sarıhan, E., Daeshian, A. M. and Moghadassi, A. S. 2007. The effect of corm size and different harvesting times on saffron (*Crocus sativus* L.) regeneration. *Acta Hort.(ISHS)*, 739: 113-117.
- Arslan, N. 1997. Safran (*Crocus sativus*L.)'da Tohumluk Materyalin Çoğaltılması. *Türkiye' de Tarım*, 1(2): 3-4.
- Basker, D. and M. Negbi. 1983. Uses of saffron (*Crocus sativus* L.). *Economic Botany*, 37: 228–236.
- DeJuan, J.A., Corcoles, H. L., Munoz R. M. and Picornell. M. R. 2009. Yield and yield components of saffron (*Crocus sativus* L.) under different cropping systems. *Ind. Crop Prod.*, 30: 212–219.
- Deo, B. 2003. Growing saffron (*Crocus sativus* L.)- the world most expensive spice. *Crop & Food Research*. Available on Internet at:www.cropcri.nz/home.
- Fernandez, J.A. 2004. Biology, biotechnology and biomedicine of saffron (*Crocus sativus* L.). *Recent Res. Devel. Plant. Sci.*, 2: 127-159.
- Gresta, F., Lombardo, G. M., Siracusa, L. and Ruberto, G. 2008. Effect of mother corm dimension and sowing time on stigma yield, daughter corms and qualitative aspects of saffron (*Crocus sativus*L.) in a Mediterranean environment. *J. Sci. Food Agric.*, 88: 1144-1150.
- Gümüşsuyu, İ. 2003. Top of the spices Saffron(*Crocus Sativus*L.). *Safranbolu Society of Service and Culture Publication, Safranbolu, Turkey, P:12*.
- Gümüşsuyu, İ. 2002. Dünyanın En Pahalı Baharatı Safran (*Crocus sativus* L.). *Safranbolu Hizmet Birliği KültürYayını*, No: 12.
- Hosseini, M., Sadeghiand, B. and Aghamiri, S. A. 2004. Influence of Foliar Fertilization on Yield of saffron (*Crocus sativus* L.). *Acta Hort.*, 650:195-200.
- İpek A., Arslan, N. and Sarıhan, E. O. 2009. Effects of different planting depth and bulb sizes on yield and yield components of saffron (*Crocus sativus* L.). *Ankara Üniversitesi, Ziraat Fakültesi, Tarım Bilimleri Dergisi*, 15(1):38-46.
- McGimpsey, J. A., Douglas, M. H. and Wallace, A. R. 1997. Evaluation of Saffron (*Crocus sativus* L.) production in New Zealand. *New Zealand J. Crop andHort. Sci.*, 25:159-168.
- Moraga, A. R., Rambla, J. L. Ahrazem, O. Granell, A. and Gomez, L. 2009. Metabolite and target transcript analyses during saffron (*Crocus sativus* L.) stigma development. *Phytochemistry*, 70: 1009–1016.
- Nehvi, F. A., Wani, S. A. Dar, S. A. Makhdoomi, M. I. Allie, B. A. and Mir, Z. A. 2007. New emerging trends on production technology of saffron (*Crocus sativus*L.). *Proc. II. IS on Saffron Bio. And Techno*.

Eds.; A. Koochehi et. al. *Acta Hort. (ISHS)*., 739: 375-381.

- Kafi, M., Rashed, M. H. Koochehi A. and Mollafilabi, A. 2002. Saffron: Production Technology and Processing. Center of Excellence for Agronomy (Special Crops). *Faculty of Agriculture, Ferdowsi University of Mashhad, Iran*.
- Vurdu, H., Çaltu, Z. and Ayan, S. 2002. Propagation techniques of Saffron plant (*Crocus sativus* L.). *J. For Fac, Gazi Univ. Kastamonu.*, 2(2): 175-187.
- Vurdu, H. and Güneş, K. 2004. SafranKırmızıAltın. *GaziÜniversitesi, KastamonuOrmanFakültesi*.
- Vurdu, H. 1993. Plants with Bulbs and Forbidden Protection. *Fidan.*, 60: 2-3.