



Antibacterial activity of the essential oils from five endemic herbs (Lamiaceae)

Behzad Hamedi*, Zeinab Mohammadi, Abdollah Ghasemi Pirbalouti

Research Center for Medicinal Plants and Ethnoveterinary, Islamic Azad University of Shahrekord Branch, 88186-34141 Shahrekord, Iran

*Email: hamedi_b123@yahoo.com

ARTICLE INFO

Type: Short Communication

Topic: Pharmacognosy

Received 4th June 2013

Accepted 26th August 2013

Key words:

- ✓ Medicinal plants
- ✓ Essential oil
- ✓ Food preservation

ABSTRACT

Background & Aim: The major aim of this study was to determine the antibacterial activity of the essential oils from five medicinal plants.

Experimental: The hydro-distillation essential oils from five plant species, including *Satureja bachtiarica*, *Zataria multiflora*, *Thymus daenensis*, *Thymus carmanicus*, and *Ziziphora tenuior* (Lamiaceae) which are endemic in Iran. *In-vitro* antibacterial test was done by agar disc diffusion and serial dilution assays.

Results & Discussion: Most of the essential oils indicated relatively antibacterial activity against the tested bacteria with the diameter of inhibition zone ranging between 7 and 33 mm.

Recommended applications/industries: The essential oils from Iranian herbs could be used as natural antimicrobial agents in food preservation.

1. Introduction

There is worldwide concern about the appearance and the rise of bacterial resistance to commonly used antibiotics. In this regard, programs for monitoring resistance have been implemented in many countries for the purpose of protecting the health of humans as well as animals (Cizman, 2003; Aarestrup, 2004). In recent years there is increasing incidence of multiple resistances in human and animal pathogenic microorganisms, largely due to the indiscriminate use of commercial antimicrobial drugs commonly employed in the treatment of infectious diseases (Westh et al., 2004). This has forced scientist to search for new antimicrobial substances from various sources like medicinal plants. Search for new antibacterial agents should be continued by the screening of many plant families (Parekh & Chanda, 2006).

Plant substances play a major role in primary health care as therapeutic remedies in many developing countries until today (Zakaria, 1991). Medicinal herbs contain physiologically active principles that over the years have been exploited in traditional medicine for the treatment of various ailments as they contain antimicrobial properties (Kelmanson et al., 2000; Srinivasan et al., 2001).

Essential oils and extracts of various species of edible and medicinal plants, herbs, and spices constitute of very potent natural biologically active agents. Use of essential oils as

antimicrobial agents in food systems may be considered as an additional intrinsic determinant to increase the safety and shelf life of foods (Nejad Ebrahimi et al., 2008). Chemical medicines, because of their harmful and irreversible effects on people, are slowly being replaced by active substances of plants (Banerjee, 2002).

Iranian medicinal plants, including *Satureja bachtiarica*, *Zataria multiflora*, *Thymus daenensis*, *Thymus carmanicus*, and *Ziziphora tenuior* have been utilized as traditional medicines by the indigenous people of Chaharmahal va Bakhtiari in Iran (Ghasemi Pirbalouti, 2009). The aim of the present study is to reveal the antibacterial properties of listed medicinal plants through *in vitro* investigation.

2. Materials and Methods

2.1. Plant material

Plants were selected for this study based on their medicinal usages in Iran. Their identity was confirmed and voucher specimens were deposited at the Research Center for Medicinal and Aromatic Plants, Islamic Azad University, Shahrekord Branch, Iran.

2.2. Essential oil extraction

The aerial parts of plants were powdered (200 g) and subjected to hydro-distillation (2000 ml distilled water) for 4 h using a Clevenger-type apparatus according to the method recommended in British Pharmacopoeia (British Pharmacopoeia, 1988). The oils were stored in universal bottles and refrigerated at 4 °C prior to use.

2.3. Bacterial strains

Six bacterial strains used in this study, including *Bacillus cereus*, *Streptococcus agalactiae*, *Staphylococcus aureus*, *Listeria monocytogenes*, *Salmonella typhimurium* and *Proteus vulgaris* were obtained from the culture collection at Persian Institute of Industrial and Scientific Research (Persian Type Culture Collection). Cultures of bacteria were grown on Blood agar (Merck, Germany).

2.4. Sensitivity testing

The sensitivity testing of the essential oils were determined using agar well diffusion method (Irobi et al., 1996). The MIC of the oils was also determined using a two-fold dilutions method. The bacterial isolates were first grown in nutrient broth for 18 h before use. The inoculum suspensions were standardized and then tested against the effect of the two plant oils at a concentration of 20 µg/ml each in DST medium. The plates were later incubated at 37°C ± 0.5°C for 24 h after which they were observed for zones of inhibition.

3. Results and Discussion

In the present study, five the essential oils from folklore medicinal plants have been tested for their antibacterial effect against six microbial strains were recorded (Table 1). Most of the oils indicated relatively antimicrobial activity against the tested bacteria with the diameter of inhibition zone ranging between 7 and 33 mm. The antimicrobial activities of herbal oils varied related to the test organism. Among the essential oils, *Satureja bachtiarica* and *Thymus daenensis* showed the best antimicrobial activity. On the other hand, concentrations of 8, 16, and 32 µg/ml of *Ziziphora tenuior* extracts had no effect on bacterial growth. The most active of the concentration was 250 µg/ml that inhibiting the growth of bacteria. The results obtained appeared to confirm the antibacterial potential of the plants investigated.

During the last years, the antimicrobial effect of medicinal plants is well documented all over the world. In Pakistan, the methanolic, hot water and cold water extracts of *Pistacia integerrima* (gall), *Polygonum Bistorta* (root), *Swetia charita* (stem), and *Zingiber officinale* (root) were

screened against gram positive bacteria, such as *Bacillus subtilis* (ATCC6633), *Enterococcus faecalis* (ATCC14506) and *Staphylococcus aureus* (ATCC6538), and gram negative bacteria like *Pseudomonas aeruginosa* (ATCC27853) and *Salmonella typhi* (ATCC14028) for their antibacterial efficiency. The maximum zone of inhibition of 19 mm of methanolic extract of *Zingiber officinale* was observed against *Staphylococcus aureus*, while the maximum zone of inhibition of 15 mm of cold water extract of *Zingiber officinale* against *Pseudomonas aeruginosa* and 15 mm of cold water extract of *Swetia charita* against *Bacillus subtilis* were observed (Khalid et al., 2011). The antimicrobial activities of the extracts of eight Iranian traditional plants, including *Hypericum scabrum*, *Myrtus communis*, *Pistachia atlantica*, *Arnebia euchroma*, *Salvia hydrangea*, *Satureja bachtiarica*, *Thymus daenensis* and *Kelussia odoratissima*, were investigated against *Escherichia coli* O157:H7, *Bacillus cereus*, *Listeria monocytogenes* and *Candida albicans* by agar disc diffusion and serial dilution assays (Ghasemi Pirbalouti et al., 2010). Most of the extracts showed a relatively high antimicrobial activity against all the tested bacteria and fungi (Ghasemi Pirbalouti et al., 2010). According to a report of Rasooli et al. (2006), various concentrations of essential oils from *Thymus eriocalyx* and *Thymus x-porlock* tested on agar plates and in broth tubes showed very strong anti-listeria properties. Also, they reported that *Thymus x-porlock* was a stronger bactericidal agent than *Thymus eriocalyx* oil.

Medicinal and aromatic plants with antimicrobial activities have become more interesting because some of them are part of the arsenal of modern antimicrobial drugs and many people are aware of problems associated with the over-prescription and misuse of traditional antibiotics. In this study, the antibacterial activity of endemic Iranian plants was investigated.

4. Conclusion

In the present study, we demonstrated the potent antibacterial activity of Iranian herbs oils against food-borne pathogens strains, which justifies the large use of this plant in traditional medicine. However, further research is needed to evaluate the effectiveness of these extract in food ecosystems to establish their utility as natural antimicrobial agents in food preservation and safety.

Table 1. Antibacterial activity of Iranian herbal extracts on six bacterial strain.

Bacterial strain	<i>S. bachtiarica</i>					
	8 (µg/ml)	16 (µg/ml)	32 (µg/ml)	64 (µg/ml)	128 (µg/ml)	256 (µg/ml)
<i>Bacillus cereus</i>	12 ^a	15	23	26	29	33
<i>Streptococcus agalactiae</i>	9	13	17	19	22	27
<i>Staphylococcus aureus</i>	12	13	14	17	16	19
<i>Listeria monocytogenes</i>	10	12	24	27	29	31
<i>Salmonella typhimurium</i>	17	19	20	22	22	26
<i>Proteus vulgaris</i>	10	12	13	16	19	24
<i>Z. multiflora</i>						
<i>Bacillus cereus</i>	9	11	13	18	19	23
<i>Streptococcus agalactiae</i>	10	11	11	17	22	27
<i>Staphylococcus aureus</i>	- ^b	-	13	16	18	19
<i>Listeria monocytogenes</i>	-	-	10	18	23	25
<i>Salmonella typhimurium</i>	-	12	13	16	23	26
<i>Proteus vulgaris</i>	12	12	12	15	23	25
<i>T. daenensis</i>						
<i>Bacillus cereus</i>	13	18	19	21	25	27
<i>Streptococcus agalactiae</i>	14	16	18	21	24	27
<i>Staphylococcus aureus</i>	11	14	16	19	23	27
<i>Listeria monocytogenes</i>	13	17	19	23	25	26
<i>Salmonella typhimurium</i>	11	15	18	23	25	29
<i>Proteus vulgaris</i>	15	18	23	25	29	33
<i>T. carmanicus</i>						
<i>Bacillus cereus</i>	11	17	19	22	24	28
<i>Streptococcus agalactiae</i>	17	19	21	24	26	27
<i>Staphylococcus aureus</i>	16	19	21	24	26	29
<i>Listeria monocytogenes</i>	15	21	24	25	28	30
<i>Salmonella typhimurium</i>	10	15	17	18	22	25
<i>Proteus vulgaris</i>	15	14	17	27	28	30
<i>Z. tenuior</i>						
<i>Bacillus cereus</i>	-	-	-	-	7	17
<i>Streptococcus agalactiae</i>	-	-	-	10	13	16
<i>Staphylococcus aureus</i>	-	-	-	-	10	15
<i>Listeria monocytogenes</i>	-	-	-	7	12	14
<i>Salmonella typhimurium</i>	-	11	12	12	13	15
<i>Proteus vulgaris</i>	-	-	-	-	16	16

^a Zone of inhibition (mm)

^b No inhibition zone

5. References

Aarestrup, F. M., 2004. Monitoring of antimicrobial resistance among food animals: principles and limitations. *Journal of Veterinary Medicine, B, Infectious Diseases and Veterinary Public Health.*, 51: 380–388.

Banerjee, M., 2002. Public policy on Ayurveda. *Economic and Political Weekly.*, 37: 1136-1146.

British Pharmacopoeia, 1988. (Vol. 2, pp. 137–138). London: HMSO.

Cizman, M., 2003. The use and resistance to antibiotics in the community. *International Journal of Antimicrobial Agents.*, 21(4): 297-307.

Ghasemi Pirbalouti, A., 2009. Medicinal plants used in Chaharmahal and Bakhtyari districts, Iran. *Herba Polonica.*, 55: 69-75.

Ghasemi Pirbalouti, A., Jahanbazi, P., Enteshari, S., Malekpoor, F. and Hamedi, B., 2010. Antimicrobial activity of some Iranian medicinal plants. *Archive Biological Science, Belgrade.*, 62 (3): 633-642.

Irobi, O. N., Moo-Young, M., Daramola, S. O., 1996. *International Journal of Pharmacognosy.*, 34(2): 87-90.

Kelmanson, J. E., Jager, A. K. and Van Staden, J. 2000. Zulu medicinal plants with antibacterial activity. *Journal of Ethnopharmacology.*, 69: 241-246.

Khalid, A., Waseem, A., Saadullah, M., Rehman, U., Khiljee, S., Sethi, A., Hassan Bin Asad, M., Rasool, F., Waqas, M. K. and Murtaza, G., 2011. Antibacterial activity analysis of extracts of various plants against gram - positive and -negative bacteria. *African Journal of Pharmacy and Pharmacology.*, 5 (7): 887-893.

Nejad Ebrahimi, S., Hadian, J., Mirjalili, M. H., Sonboli, A., Yousefzadi, M., 2008. Essential oil composition and antibacterial activity of *Thymus caramanicus* at different phenological stages. *Food Chemistry.*, 110: 927-931.

Parekh, J. and Chanda, S., 2006. In vitro antimicrobial activities of extract of *Launaea procumbens* Roxb. (Labiatae), *Vitis vinifera* (Vitaceae) and *Cyperus rotundus* (Cyperaceae). *African Journal of Biomedical Research.*, 9: 89-93.

Rasooli, I., Rezaei, M. B. and Allameh, A., 2006. Ultrastructural studies on antimicrobial efficacy of thyme

- essential oils on *Listeria monocytogenes*. *International Journal of Infection Disease.*, 10: 236-241.
- Srinivasan, D., Sangeetha, N., Suresh, T. and Perumalsamy, P. L., 2001. Antimicrobial activity of certain Indian medicinal plants used in folkloric medicine. *Journal of Ethnopharmacology.*, 74: 217-220.
- Westh, H., Zinn, C. S., Rosdahl, V. T., 2004. An international multicenter study of antimicrobial consumption and resistance in *Staphylococcus aureus* isolates from 15 hospitals in 14 countries. *Microbial Drug Resistance.*, 10: 169-176.
- Zakaria, M., 1991. Isolation and characterization of active compounds from medicinal plants. *Asia Pacific Journal of Pharmacology.*, 6: 15-20.