



Effect of ellagic acid on oxidative stress duo to brain ischemia/hypoperfusion in male rat

Maryam Rafieirad^{1*}, **Khadijeh Ghasemzadeh-Dehkordi**²

1. Department of Biology, Izeh Branch, Islamic Azad University, Izeh, Iran

2. Department of Biology, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran

* Corresponding author, *Email: Rafieirad.m@gmail.com, Rafieirad.m@izehiau.ac.ir

ARTICLE INFO

Type: Original Research

Topic: Physiology & Pharmacology

Received 17th May 2014

Accepted 27th June 2014

Key words:

- ✓ Brain ischemia
- ✓ Ellagic acid
- ✓ Malondialdehyde
- ✓ Oxidative stress
- ✓ Thiol

ABSTRACT

Background & Aim: Free radicals are produced in ischemic processes. Nerve damage caused by free radicals may play a role in neurological diseases and antioxidants are protective activity. Ellagic acid is a polyphenol compound with antioxidant properties which is found in fruits like pomegranate, blackberry, and all types of mulberry. This study aimed to evaluate the effect of 14 days of oral administration of ellagic acid (50 mg/kg) on brain oxidative stress indices after permanent bilateral common carotid artery occlusion or ischemia/hypoperfusion in male adult rats.

Experimental: A number of 30 rats were divided into test and control groups. To make animal model of permanent cerebral hypoperfusion/ischemia, carotid arteries were ligatured as upper and lower and cut bilaterally. Rat brain tissue was extracted to separate hippocampus and measure malondialdehyde, and thiol (-Sh) groups.

Results & Discussion: Our results indicated that malondialdehyde and thiol in ischemia group has significantly increased ($p < 0.001$) than to control group. In addition, the rate of malondialdehyde and thiol in ischemia group receiving the ellagic acid has significantly decreased ($p < 0.001$), than to ischemia group.

Recommended applications/industries: Ellagic acid possibly with powerful antioxidant properties, can improve the effect of ischemia such as production of free radicals.

1. Introduction

Ischemia means reducing cerebral blood flow (CBF) which damages the whole or part of brain (Rafiei Rad *et al.*, 2011). 2-VO (Two Vessel Occlusion) model is an experimental model for chronic brain hypoperfusion, in which permanent occlusion of both common carotid arteries and their cutting occur (in rats) (Accardo-Palumbo *et al.*, 2010). In normal

conditions, there is a balance between production and elimination of free radicals and an imbalance in these processes leads to oxidative stress and incidence of pathological changes in multiple cellular macromolecules (Farkas *et al.*, 2007). As a result, oxidative stress can alter neurotransmission, neuronal function and whole brain activity (Rammal *et al.*, 2007). One of factors causing oxidative stress is reactive oxygen species (ROS) which is generated

within brain tissue during ischemia, and plays a role in the development of cerebral damages (Gil *et al.*, 2000). ROS are highly reactive and attack lipids, proteins, and nucleic acids, which eventuates in tissue injury and cell death (Gil *et al.*, 2000). Interestingly, oxidative stress state was recently linked to other behavioral disorders, such as aggressive behavior and depression, and also to deterioration of short-term spatial memory (Bouayed *et al.*, 2009a; Bouayed *et al.*, 2009b), highlighting that oxidative stress disturbances could be implicated in the pathophysiology of conditions that are more specific for the nervous system impairment (Bouayed *et al.*, 2009b).

In recent years, it has been well shown that consuming fruits and vegetables containing natural antioxidants prevents many diseases such as cardiac complications and even different cancers (Casanova *et al.*, 2008). Ellagic acid (8, 7, 3, 2- tetra-hydroxy-chromno (3, 4, 5-CDA) chromno-10, 5-dion with molecular weight of 302 g/mole) is a bioactive compound with many pharmaceutical and industrial applications (Ascacio-Valdes *et al.*, 2010). This molecule has different properties such as antioxidant ones (Ascacio-Valdes *et al.*, 2010). The studies conducted on ellagic acid in cancer cells have shown apoptosis induction, cellular death and stopping continual growth of tumor (Ito *et al.*, 1999). Results of studies suggest that ellagic acid provides effects of useful neural protection against oxidative damage in diabetic rats (Uzar *et al.*, 2012).

In this research, effects of ellagic acid on oxidative stress in ischemia hypoperfusion model were investigated.

2. Materials and Methods

2.1. Animals and experimental groups

The present research was done using adult male Wistar rats reproduced in Reproduction and Care Center, Animal House of Ahwaz University of Medical Sciences, in weight range of 200-250 g. The animals were being kept inside standard cages in quartet groups in Animal Care Center of Islamic Azad University, Izeh Branch, Khuzestan, Iran. They were kept in standard conditions at 20±2 °C and 12-h light-dark cycle (lighting started at 7 a.m.) with enough access to compressed food of Dam Pars Tehran Company,

Chavdaneh Company, Shahreza, Isfahan, and purified tap water of Izeh City. To facilitate the work and adapt to peripheral conditions and the tester, the animals were petted for some minutes on a daily basis in advance. They were randomly classified into the following groups:

1. Healthy group without ischemic induction and medicine intake (control);
2. Ischemic hypoperfusion group without medicine intake (ischemia);
3. Ischemic hypoperfusion group receiving 50 mg/kg of ellagic acid every day for 14 days using intragastric administration or gavage (ischemia + 50 mg/kg ellagic acid) (Girish *et al.*, 2013; Rafieirad *et al.*, 2014a)

2.2. Brain ischemia/hypoperfusion Surgery Method

After completely anesthetizing the animals using Ketamin / xylazine (intra protaneal injection of 10.90 mg / kg of body weight), left and right common carotid arteries were opened by longitudinal incision of skin below their neck and pushing muscles aside. After making two knots using surgical suture silk on top and at bottom of each artery, middle of both knots was cut by a pair of scissors. Then, places of the surgery were sutured and then the animals were treated to recover from the surgery. After the animals became conscious after one week, a similar surgery was performed on the other side (Rafieirad *et al.*, 2014). Afterward, the animals of both groups were treated for two weeks with one of different doses of ellagic acid using gavage method.

2.3. Preparation of ellagic acid

Ellagic acid from Sigma-Aldrich (Steinheim, Germany) was solved in normal saline. It was administered orally by gavage for 14 days (Rafieirad *et al.*, 2014a).

2.4. Brain Sample Collection and Biochemical Assays

At the end of experiments, the animals were decapitated and the hippocampus was removed quickly and rinsed with saline. The tissues were immediately homogenized in cold KCl solution (1.5%) to obtain 10% homogenate suspension used for measuring thiobarbituric acid reactive substances (TBARS) value, expressed as malondialdehyde equivalents (MDA) (Sarkaki *et al.*, 2013; Rafieirad *et al.*, 2014b) (Fig 1.).




Fig 1. The Standard Curve of MDA was constructed Over the Concentration Range of 0–10 μ M

2.6. Statistical analysis

Data of this research were analyzed using one way ANOVA by SPSS_{ver 20.0} software. Means were comprised LSD test methods with $p < 0.05$ and then they presented as mean \pm SEM.

3. Results and discussion

In this study, in which effect of ellagic acid was studied on antioxidant status such as lipid peroxidation and thiol group (-Sh) in animal models of ischemia, malondialdehyde (MDA) significantly increased in hippocampus ($p \leq 0.001$) in the ischemic group compared with the control. The observations showed that malondialdehyde in hippocampus tissue in ischemia group treated with (50 mg/kg) of ellagic acid significantly reduced ($p \leq 0.001$) compared with the ischemia group (Fig 2.). On the other thiol in ischemia group has significantly increased ($p \leq 0.001$) than in control group and rate of thiol in ischemia group receiving ellagic acid has significantly decreased ($p \leq 0.001$) (Fig 3.).

As can be seen in figure our results indicated that ellagic acid could reduce oxidative stress factors that caused by ischemia and these favorable effects were significant. Cerebral ischemia caused increase of Reaction Oxygen Species (ROS) and finally activation of the pathways leading to cellular death in vulnerable areas of brain (Puyal *et al.*, 2013). Neurons have the highest sensitivity to ischemia in CA1 area of hippocampus (Eichenbaum, 2004). Previous researches have shown that obstruction of both common carotid

arteries for 30 min caused extensive cellular death in CA1 area of hippocampus so that significant reduction was found in mean number of CA1 area in ischemic group compared with other groups and significant changes were found in behavioral tests of the animals (Hazelton *et al.*, 2010). Oxidative stress not only increases production of free radicals, but also reduces cellular antioxidant mechanisms.




Fig 2. Mean \pm SD of MDA between Control Group, Ischemia and Ischemic Group orally receiving 50 mg/kg ellagic acid for 14 Days.




Fig 3. Mean \pm SD of total thiol between Control Group, Ischemia and Ischemic Group orally receiving 50 mg/kg ellagic acid for 14 Days

Evaluation of oxidative stress is done using different indicators, one of which is the end products of lipid peroxidation (MDA). MDA with thiobarbituric acid (TBA) creates a red fluorescent derivative that is measured by spectrophotometrically (Acikgoz *et al.*, 1998). Philip *et al.* (2003) found that after 40 minutes of ischemia, the level of total protein S - thiolation significantly increased in kidney tissue and remained significantly higher than control after 5 min

reperfusion. Low molecular weight thiols such as glutathione can act as a protection against the irreversible loss of biological activity of macromolecules. Glutathione is synthesized in two reactions, gamma-glutamyl cysteine synthetase by successive ATP-dependent (γ -GCS), the rate-limiting enzyme, glutathione synthetase (Eaton *et al.*, 2003). GSH concentrations is inhibited by a feedback mechanism of γ -GCS activity (Lu, 2000). Results have shown that increasing glutathione levels in old mice significantly improved by grape seed extract, as to justify that flavonoids in grape seed extract produces the rate-limiting enzyme in the synthesis of glutathione (gamma-glutamyl cysteine synthetase) (Balu *et al.*, 2005). Recent studies have focused on the role of antioxidants in the treatment of nervous system disorders and possibly ischemia, which is based on the assumption that if oxidative stress occurs when there is an imbalance between free radical production and antioxidant availability, administration of supplemental antioxidants may be used to sweep radicals and change disease development, progression or both. Recently, health food or nutritional supplements have attracted much of public attention (Motamedi *et al.*, 2010). Ellagic acid is one of the most important compounds available in pomegranate and phenol structure and nature of this compound causes its strong antioxidant activity (Sopala & Danysz, 2001).

Results a study by Spencer *et al.* (2011) indicated that ellagic acid prevented oxidative damage of DNA due to copper transporters and catecholamine; as a result, protective role of ellagic acid has been suggested for preventing ROS, lipid peroxidation and breakage of DNA string (Spencer *et al.*, 2011). Recent studies have investigated ability of cytotoxic activity and antigrowth factor of ellagic acid in human cells, lung cancer, colorectal cancer, breast cancer and prostate cancer and shown that 1 to 100 M/lit inhibits expansion of cell cancers. In addition, it has been demonstrated that application of ellagic acid probably leads to apoptosis in cancer cells by inhibiting the factors that cause metastasis (Spencer *et al.*, 2011). The ellagic acid has been recognized as antimicrobial and antiviral factor and it has been also mentioned that ellagic acid is able to inhibit growth of pathogens in humans; this action is probably performed by being coupled with protein in wall of bacteria (Huetz *et al.*, 2005). In addition, ellagic acid has some inflammatory effects and causes reduced oxidative stress (Chao *et al.*, 2009). According to the

results of this study, confirm previous reports functional disorders after brain ischemia that is caused by damage on brain cells is due to oxidant production.

4. Conclusion

Therefore, ellagic acid with antioxidant ability to remove oxidant substances from specific and important areas of the brain can be this compensatory effect on oxidative stress in animal model of hypoperfusion ischemia.

6. References

- Accardo-Palumbo, A., D'Amelio, L., Pileri, D., D'Arpa, N., Mogavero, R., Amato, G., Cataldo, V., Napoli, B., Ciccia, F., Lombardo, C. and Conte, F., 2010. Reduction of plasma granzyme A correlates with severity of sepsis in burn patients. *Journal of the International Society for Burn Injuries.*, 36: 811-818.
- Acikgoz, O., Gonenc, S., Kayatekin, B. M., Uysal, N., Pekcetin, C. Semin, I. and Gure, A., 1998. Methamphetamine causes lipid peroxidation and an increase in superoxide dismutase activity in the rat striatum. *Brain Research.*, 813: 200-202.
- Allahbakhshi, E., Rafieirad, M. and Zangeneh Nezhad, Z., 2014. Neuroprotective effects of oral ellagic acid on locomotor activity and anxiety-induced by ischemia/hypoperfusion in rat., *Advances in Environmental Biology.*, 8: 83-88.
- Ascacio-Valdes J.A., Aguilera-Carbo, A., Martínez-Hernandez, J.J., Rodríguez-Herrera, R. and CN, A. 2010. *Euphorbia antisiphilitica* residues as a new source of ellagic acid. *Chemical Papers.*, 64: 528-532.
- Balu, M., Sangeetha, P., Murali, G. and Panneerselvam, C., 2005. Age-related oxidative protein damages in central nervous system of rats: modulatory role of grape seed extract. *International Journal of Developmental Neuroscience.*, 23: 501-507.
- Balu, M., Sangeetha, P., Murali, G. and Panneerselvam, C., 2005. Age-related oxidative protein damages in central nervous system of rats: modulatory role of grape seed extract. *International Journal of Developmental Neuroscience.*, 23: 501-507.

- Bouayed, J., Desor, F. and Soulimani, R., 2009. Subacute oral exposure to benzo [*alpha*] pyrene (B[*alpha*]P) increases aggressiveness and affects consummatory aspects of sexual behavior in male mice. *Journal of Hazardous Materials.*, 169: 581-585.
- Bouayed, J., Rammal, H. and Soulimani, R. 2009b. Oxidative stress and anxiety: relationship and cellular pathways. *Oxidative Medicine and Cellular Longevity.*, 2: 63-67.
- Casanova, E., Garcia-Mina, J. M. and Calvo, M. I., 2008. Antioxidant and antifungal activity of *Verbena officinalis* L. leaves. *Plant Foods for Human Nutrition.*, 63: 93-97.
- Chao, P. C., Hsu C. C. and Yin, M. C., 2009. Anti-inflammatory and anti-coagulatory activities of caffeic acid and ellagic acid in cardiac tissue of diabetic mice. *Nutrition & Metabolism (Lond).*, 6: 33.
- Eaton, P., Jones, M. E., McGregor, E., Dunn, M. J., Leeds, N., Byers, H. L., Leung, K. Y., Ward, M. A., Pratt J. R. and Shattock, M. J., 2003. Reversible cysteine-targeted oxidation of proteins during renal oxidative stress. *Journal of the American Society of Nephrology.*, 14: S290-296.
- Eichenbaum, H., 2004. Hippocampus: cognitive processes and neural representations that underlie declarative memory. *Neuron.*, 44: 109-120.
- Farkas, E., Luiten P. G. and Bari, F., 2007. Permanent, bilateral common carotid artery occlusion in the rat: a model for chronic cerebral hypoperfusion-related neurodegenerative diseases. *Brain Research Review.*, 54: 162-180.
- Gil, M. I., Tomas-Barberan, F. A., Hess-Pierce, B., Holcroft, D. M. and Kader, A. A., 2000. Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing. *Journal of Agricultural and Food Chemistry.*, 48: 4581-4589.
- Girish, C., Raj, V., Arya, J. and Balakrishnan, S., 2013. Involvement of the GABAergic system in the anxiolytic-like effect of the flavonoid ellagic acid in mice. *European Journal of Pharmacology.*, 710: 49-58.
- Hazelton, J. L., Balan, I., Elmer, G. I., Kristian, T., Rosenthal, R. E., Krause, G., Sanderson, T. H. and Fiskum, G., 2010. Hyperoxic reperfusion after global cerebral ischemia promotes inflammation and long-term hippocampal neuronal death. *Journal Neurotrauma.*, 27: 753-762.
- Huetz, P., Mavaddat, N. and Mavri, J., 2005. Reaction between ellagic acid and an ultimate carcinogen. *Journal of Chemical Information Modeling.*, 45: 1564-1570.
- Ito, H., Miyake, M., Nishitani, E., Mori, K., Hatano, T., Okuda, T., Konoshima, T., Takasaki, M., Kozuka, M. Mukainaka, T. Tokuda, H. Nishino, H. and Yoshida, T., 1999. Anti-tumor promoting activity of polyphenols from *Cowania mexicana* and *Coleogyne ramosissima*. *Cancer Letterer.*, 143: 5-13.
- Lu, S. C. 2000. Regulation of glutathione synthesis. *Current Topics in Cellular Regulation.*, 36: 95-116.
- Mansouri, M. T., Farbood, Y., Sameri, A. Sarkaki, M. J. Naghizadeh, B. and Rafeirad, M., 2013. Neuroprotective effects of oral gallic acid against oxidative stress induced by 6-hydroxydopamine in rats. *Food Chemistry.*, 138: 1028-1033.
- Motamedi, F., Rafiei Rad, M., Sarkaki, A., Hoseini, E. and Farbood, Y., 2011. The Effect of Grape seed extract on lipid peroxidation duo to ischemia/hypoperfusion in male rat striatum. *Journal of Animal Biology.*, 3: 37-44.
- Motamedi, H., Darabpour, E., Gholipour, M. and Seyyed Nejad, S. M., 2010. In vitro assay for the anti-*Brucella* activity of medicinal plants against tetracycline-resistant *Brucella melitensis*. *Journal of Zhejiang University Science B.*, 1: 506-511.
- Puyal, J., Ginet, V. and Clarke, P. G. 2013. Multiple interacting cell death mechanisms in the mediation of excitotoxicity and ischemic brain damage: a challenge for neuroprotection. *Progress in Neurobiology.*, 105: 24-48.
- Rafiei Rad, M., Sarkaki, A., Hoseini, E., Farbood, Y. and Mansouri, S.M.T., 2011: The effect of grape seed extract on lipid peroxidation duo to ischemia/hypoperfusion in male rat striatum. *Journal of Animal Biology.*, 3: 37-44.
- Rafieirad, M., Zangeneh Nezhad, Z. and Allahbakhshi, E., 2014a. Neuroprotective effects of oral ellagic acid on locomotor activity and anxiety-induced by ischemia/hypoperfusion in rat. *Advances in Environmental Biology.*, 8: 83-88.
- Rafieirad, M., Zangeneh Nezhad, Z. and Eydipour, Z., 2014b. Effects of oral *Ferulago angulata* hydroalcoholic extract on lipid peroxidation induced by ischemia/hypoperfusin in the

hippocampus rat. *International Journal of Biology, Pharmacy and Allied Sciences*, 3, 157-164.

- Rammal, H., Bouayed J. and Soulimani, R. 2007. A direct relationship between aggressive behavior in the resident/intruder test and cell oxidative status in adult male mice. *European Journal of Pharmacology.*, 627: 173-176.
- Sarkaki, A., Rezaiei, M., Gharib Naseri, M. and Rafieirad, M., 2013. Improving active and passive avoidance memories deficits due to permanent cerebral ischemia by pomegranate seed extract in female rats. *Malaysian Journal of Medical Sciences.*, 20: 25-34.
- Sopala, M. and Danysz, W., 2001. Chronic cerebral hypoperfusion in the rat enhances age-related deficits in spatial memory. *Journal of Neural Transmission.*, 108: 1445-1456.
- Spencer, W. A., Jeyabalan, J., Kichambre, S. and Gupta, R. C., 2011. Oxidatively generated DNA damage after Cu (II) catalysis of dopamine and related catecholamine neurotransmitters and neurotoxins: Role of reactive oxygen species. *Free Radical Biology and Medicine.*, 50: 139-147.
- Uzar, E., Alp, H., Cevik, M. U., Firat, U., Evliyaoglu, O., Tufek, A. and Altun, Y. 2012. Ellagic acid attenuates oxidative stress on brain and sciatic nerve and improves histopathology of brain in streptozotocin-induced diabetic rats. *Neurol Science.*, 33: 567-574.