



## Chemical composition and antioxidant activity of *Acorus calamus* L. accessions from different altitudes of Uttarakhand Himalayas

Archana Parki, Pinky Chaubey, Om Prakash\*, Ravindra Kumar and Anil K. Pant

Department of Chemistry, College of Basic Sciences and Humanities, G.B. Pant University of Agriculture and Technology, Pantnagar, U.S. Nagar-263145, Uttarakhand, India;

\*Email: [oporgchem@gmail.com](mailto:oporgchem@gmail.com); [parki.archana15@gmail.com](mailto:parki.archana15@gmail.com)

### ARTICLE INFO

**Type:** Original Research

**Topic:** Medicinal Plants

**Received** February 24<sup>th</sup> 2018

**Accepted** May 26<sup>th</sup> 2019

### Key words:

- ✓ *Acorus calamus*
- ✓ Antioxidant activity
- ✓ Total phenols
- ✓ Asarone
- ✓ Shyobunone

### ABSTRACT

**Background & Aim:** *Acorus calamus* L. (Araceae) is an herb traditionally used in Indian and Chinese system of medicine. It is a perennial herb with long, cylindrical scented rhizomes which are creeping and extensively branched with the thickness up to 2.5 cm. The rhizomes are whitish internally and brownish externally in appearance. It is vernacularly known as Bach in India. The present investigation reveals the phytochemical screening and antioxidant activity of the rhizome and leaves methanolic extracts of *A. calamus* from three different altitudes of Uttarakhand.

**Experimental:** The extracts compositions of *Acorus calamus* rhizomes and leaves collected from three different altitudes of Uttarakhand Himalayas, India. The yield of extracts ranged from 0.3- 4.8% w/v, were analyzed by GC/MS and in-vitro antioxidant assay were done by different methods.

**Results:** Over 65 contributing 80.90-90.55 % in different extracts among all the accessions were identified. The major compounds identified were  $\beta$ -asarone (44.9-51.9%), shyobunone (1.1%- 5.3%), Z-methyl isoeugenol (0.1%-2.4%), linoleic acid (6.4%-18.9%),  $\alpha$ -asarone(0.1%-4.6%) and Z-isoelemicin (2.2%-15.8%). The amount of phenolics in rhizome extracts, ranged from 4.10 mg-4.80 mg GAE/g respectively, whereas in leaf extracts the amounts were 2.40-3.26mg GAE/g respectively. All the extracts exhibited good in vitro antioxidant activity with the IC<sub>50</sub> values ranging from 0.3- 4.8 % w/v determined by different methods compared to standard antioxidant.

**Recommended applications/industries:** Based on above observations it can be inferred that the herb may be a good source of bioactive compounds and can work as an antioxidant to prevent the oxidative deteriorative activity of food materials beside generation of database for its scientific and judicious in-situ exploitation.

### 1. Introduction

Sweet flag, *Acorus calamus* L., vernacularly known as Bach in India is, a semi-aquatic herbaceous plant growing in temperate to sub temperate regions. It is a perennial herb with long, cylindrical rhizomes which are creeping and extensively branched with the

thickness up to 2.5 cm. The rhizomes are whitish internally and brownish externally in appearance. The rhizomes and leaves of this plant have been used in the Indian systems of traditional medicine for hundreds of years. The leaves and rhizomes essential oils of this herb have been reported to possess biological activities

like antispasmodic, carminative and also used for treatment of epilepsy, mental ailments, chronic diarrhea, dysentery, bronchial catarrh, intermittent and tumors (Avadhani *et al.*, 2016). It has been reported that the extracts are used in the traditional *Chinese* prescriptions and its beneficial effects on memory disorders and on learning performance. Lipid peroxidase content and anti-aging effect in senescence have also been reported (Mehrotra *et al.*, 2003). The plant has also been reported to possess insecticidal, antifungal, antimicrobial, moth, mosquitos, lice repellent, and as a toxicant for flying insects, ectoparasites and insects in stored products (Lewis and Elvin-Lewis, 2003; Phongpaichit *et al.*, 2005). In view of its traditional uses as medicinal herb in indigenous system, the present communication deals with chemical composition and antioxidant assay. The chemical composition along with biological activities of essential oil of this herb has already been reported by our group (Joshi *et al.*, 2012; Kumar *et al.*, 2009).

## 2. Materials and Methods

### 2.1. Plant material

The samples were selected from different altitudes of Uttarakhand Himalayas. The plants were collected from Pithoragarh, District- Pithoragarh (altitude 1514m), Bhimtal, District- Nainital (altitude 1370m) and Pantnagar District Udham Singh Nagar (altitude 344m) from their natural habitat. The rhizomes and leaves were shade dried and were powdered. The powdered material (100gm) was subjected for extraction in methanol by cold percolation method for seven days with constant shaking in between. The yield of extracts ranged from 0.3- 4.8% w/v.

### 2.2. GC-MS analysis

The GC-MS data were analysed by using GCMS-QP2010 Plus with following experimental conditions: Initial temperature 60°C with RAMP of 3°C/ min, final temperature 210°C, final hold time 10 min, carrier gas He, flow rate 1mL/ min. Column, silica DB-5, capillary(30m× 0.25mm,0.25 µm). MS were recorded under electron ionization (EI) condition (70 eV) with split mode of 40:1. The compounds were identified by matching their mass spectra with those recorded in NISTMS Wiley Library and comparing the data in literature (Adams, 2007).

### 2.3. Antioxidant activity

#### 2.3.1. DPPH radical scavenging activity

It is a quick method to study the scavenging ability of the antioxidants. A standard protocol was followed for the same (Burits and Bucar, 2000). In brief different amounts of the tested sample (20-100 µg/mL) were added to 5 mL of a 0.004% methanol solution of DPPH. Finally the absorbance was read against a blank at 517 nm after 30 min of incubation at room temperature. All the observations were taken as triplicate. BHT (butylated hydroxyl toluene) and catechin were used as the standard antioxidant. Percent DPPH free radical Inhibition (IC%) was calculated by using the equation,  $IC \% = (A_0 - A_t/A_0) \times 100$ , where  $A_0$  and  $A_t$  are the absorbance values of the control and test sample respectively.

#### 2.3.2. Reducing power activity

The reducing power of extracts was determined by following reported methods (Jeena *et al.*, 2016). Varying concentrations of tested sample (20-100 µg/mL) were mixed with 2.5 mL of phosphate buffer (200 mM, pH= 6.6) and 2.5 mL of 1% potassium ferricyanide,  $K_3[FeCN_6]$ . The mixtures were incubated for 20 min. at 50°C. After incubation, 2.5 mL of  $CCl_3COOH$  was added to the mixtures, followed by centrifugation at 650 rpm for 10 min. The supernatant (1 mL) was mixed with 5mL distilled water and 1 mL of 0.1%  $FeCl_3$ . The absorbance of the resultant solutions was measured at 700 nm. BHT was taken as the standard. The reducing power of samples was calculated by the following formula, Reducing power  $\% = (A_0 - A_t/A_0) \times 100$ , where  $A_0$  and  $A_t$  are the absorbance values of the control and test sample respectively.

#### 2.3.3. Metal chelating activity

The chelation of  $Fe^{2+}$  by extracts were evaluated by using the method developed earlier (Kunwar *et al.*, 2013). In brief 0.1 mL of 2mM  $FeCl_2 \cdot 4H_2O$ , 0.2mL of 5mM ferrozine and 4.7 mL of methanol was added to different concentrations of tested sample (20-100 µg/mL). The solutions were mixed and allowed to react for 10 min. The absorbance at 562 nm was measured. EDTA and citric acid was taken as the standards.

### 2.3.4. Total phenolics assay

In order to correlate the antioxidant activity of extracts the total phenolic content were estimated quantitatively by using the reported method (Shetty *et al.*, 1995). Briefly, 1 mL of the sample extract was transferred into a test tube and mixed with 1 mL of 80% methanol and 8 mL of distilled water. To each sample 0.5 mL of 1 N Folin-Ciocalteu reagent (FCR) was added and mixed. After 5 min., 1 mL of saturated Na<sub>2</sub>CO<sub>3</sub> solution was added to the reaction mixture and allowed to stand for 60 min. The absorbance was read at 650 nm using a Thermo Scientific UV spectrophotometer. The calibration curve was established using various concentrations of catechol and results were expressed as mg of catechol per gram of sample in dried weight (dw).

### 2.4. Statistical analysis

The experimental data were obtained by performing the experiments in triplicates. The data were analysed by Tukey's test in conjunction with an ANOVA (post-hoc) analysis with the help of SPSS 16 version.

## 3. Results and discussion

### 3.1. Phytochemical analysis of extracts

The constituents of leaves and rhizomes methanolic extracts from three populations viz., ACPGLME = *A. calamus* pithoragarh leaves methanolic extract, ACPNLME = *A. calamus* Pantnagar leaves methanolic extract, ACBTLME = *A. calamus* Bhimtal leaves methanolic extract, ACPGRME = *A. calamus* Pithoragarh rhizomes methanolic extract, ACPNRME = *A. calamus* Pantnagar rhizomes methanolic extract, ACBTRME = *A. calamus* Bhimtal rhizomes methanolic extract, were analyzed and compared using GC-MS. The analysis led to the identification of over 62 constituents comprising 80.70-90.07 % of total compositions. Pursual of table 1 shows the relative percentage of the identified components in the methanolic extracts. Phenyl propanoids contributed the major part of the total compositions, being dominated by  $\beta$ -asarone (44.9-51.9%). The other constituents identified within this class were  $\alpha$ -asarone (0.1-4.6 %), Z- isoelemicin (2.2- 15.8%), stigmast-4-en-3 one (0.3-

1.3 %), *trans* methyl isoeugenol (0.1-2.4 %), *cis* methyl isoeugenol (0.-2.0 %) and  $\gamma$ -asarone (0.5-0.7%) Besides phenyl propanoids;  $\beta$ - gurjunene (0.3-0.9%, cuparene (2.4-3%), shyobunone (1.1- 5.3%), phytol (2.3-10.6%), linoleic acid (6.4- 18.9%), 6-epishyobunone (0.8- 2.0%) were the major sesquiterpenoids identified.

$\beta$ -asarone (70.1%),  $\alpha$ -asarone (6.11%),  $\gamma$ -asarone (1.0%), methyl linoleate (0.73%), isocalamendiol (0.50%), methyl palmitate (0.39%), and methyl isoeugenol (0.26%) from hexane extracts of rhizomes have been reported in previous study (Adfa, 2015). 2,4,5-trimethoxy benzaldehyde with its activity against insects has been reported from Bangladesh (Hossain *et al.*, 2008). Saponins, resins, alkaloids, steroids, tannins, phenolics, flavonoids, proanthocyanidins, glycosides, diterpenes and triterpenes have also been reported from *A. calamus* (Barua, *et al.*, 2014; Nanda, 2014; Pandey *et al.*, 2009). The sesquiterpenes, 1 $\beta$ ,7 $\alpha$ (H)-cadinane-4 $\alpha$ ,6 $\alpha$ ,10 $\alpha$ -triol, 1 $\alpha$ ,5 $\beta$ -guaiane-10 $\alpha$ -O-ethyl-4 $\beta$ ,6 $\beta$ -diol and 6 $\beta$ ,7 $\beta$ (H)-cadinane-1 $\alpha$ ,4 $\alpha$ ,10 $\alpha$ -triol in ethanolic extracts from China have also been reported (Dong *et al.*, 2010). Shyobunone, with its insecticidal activity and linoleic acid's anti-inflammatory, acne reductive, and moisture retentive properties have also been reported (Chen *et al.*, 2015; Diezel *et al.*, 1993; Letawe *et al.*, 1998; Darmstadt *et al.*, 2002).

The constituents in various extracts are different both qualitatively and quantitatively from the results reported in the literature. The chemical diversity was also found among the extracts both from leaves and rhizomes which might be possibly because of climatic, altitudinal or edaphic factors. To the best of our knowledge and search of literature it is the first report on GC – MS analysis of methanolic extracts from rhizome and leaves of *A. calamus*. In continuation to our research on *A. calamus* we have reported the essential oil composition of *A. calamus* revealing the presence of major compounds like  $\beta$ -asarone,  $\alpha$ -asarone, Z-methyl isoeugenol,  $\beta$ -caryophyllene, methyl eugenol, spathulenol,  $\alpha$ -calacorene and caryophyllene oxide etc. along with biological activities of essential oil as a whole (Joshi *et al.*, 2012; Kumar *et al.*, 2009).

**Table 1:** Chemical constituents in methanolic extracts from leaves and rhizomes of *A. calamus*

S.N.	Compound name	% Contribution in extracts						Method of identification
		ACPGR	ACPLG	ACPNR	ACPNLM	ACBTRM	ACBTL	
1	Syringol	-	-	-	0.5	-	-	M <sup>+</sup> 154, M/z: 154,139,93
2	Linalool	0.2	1.1	-	-	0.3	2.6	M <sup>+</sup> 154, M/z: 71,93,55
3	caryophyllene	-	-	-	-	-	0.2	M <sup>+</sup> 204, M/z: 93,669,133
4	β gurjunene	0.9	-	0.4	-	0.3	-	M <sup>+</sup> 204, M/z: 161,105, 41
5	trans-methyl isoeugenol	0.7	0.6	0.1	0.1	2.4	1.1	M <sup>+</sup> 178, M/z: 178,107,163
6	γ- muurolene	-	-	0.1	-	0.1	-	M <sup>+</sup> 204, M/z: 161,105,119
7	δ cadinene	-	-	-	-	0.2	-	M <sup>+</sup> 204, M/z: 161,119,105
8	β asarone	51.9	45.9	47.0	44.9	46.2	51.2	M <sup>+</sup> 208, M/z: 208,193,165
9	Cuparene	2.6	-	3.0	-	2.4	-	M <sup>+</sup> 202, M/z: 132,145,107
10	methyl hexadecanoate	-	-	-	-	0.8	-	M <sup>+</sup> 270, M/z: 74, 87, 143
11	asaronic acid	-	1.6	-	0.8	0.3	-	M <sup>+</sup> 212, M/z: 197, 137, 169
12	hexadecanoic acid	4.8	-	5.1	-	7.9	2.0	M <sup>+</sup> 256, M/z: 73,60,43
13	linoleic acid, methyl ester	1.0	-	0.6	1.2	1.5	-	M <sup>+</sup> 294, M/z: 67, 81, 41
14	α- copaene	0.1	-	.02	-	t	-	M <sup>+</sup> 204, M/z: 161, 105, 119
15	decanoic acid	-	-	-	-	t	-	M <sup>+</sup> 172, M/z: 60, 73, 129
16	bicyclogermacrene	t	-	-	-	t	-	M <sup>+</sup> 204, M/z: 121, 93, 41
17	β- elemene	0.3	-	0.2	-	t	-	M <sup>+</sup> 204, M/z: 81,93,68
18	β-farnesene	-	-	-	-	0.1	-	M <sup>+</sup> 204, M/z: 41, 69, 93
19	Shyobunone	5.3	-	3.0	-	1.4	1.1	M <sup>+</sup> 220, M/z: 150,83,81
20	γ asarone	-	0.5	-	0.7	-	0.6	M <sup>+</sup> 208, M/z: 208, 193, 69
21	γ gurjunene	-	-	-	-	-	1.5	M <sup>+</sup> 204, M/z: 81, 107, 161
22	Elemol	-	-	-	-	-	0.3	M <sup>+</sup> 222, M/z: 59,93,161
23	hydroxycineol	-	-	-	-	0.2	-	M <sup>+</sup> 170, M/z: 43, 108, 126
24	z- isoelemicin	-	15.8	-	2.2	-	9.0	M <sup>+</sup> 208, M/z: 208,193,124
25	α-calacorene	-	-	-	-	0.07	-	M <sup>+</sup> 200, M/z: 157,142,200
26	4-propylbiphenyl-4-carboxylic acid	-	-	-	-	0.4	-	M <sup>+</sup> 240, M/z: 211, 240, 165
27	Spathulenol	0.4	-	0.5	-	0.5	-	M <sup>+</sup> 220, M/z: 43, 41, 91
28	isocalamendiol	0.8	-	0.8	-	0.7	-	M <sup>+</sup> 238, M/z: 43, 111
29	9 octadecenoic acid (z)	-	-	-	-	0.1	-	M <sup>+</sup> 282, M/z: 43, 73, 57
30	2,4,6 trimethoxy acetophenone	-	0.8	-	0.4	.07	-	M <sup>+</sup> 210, M/z: 195, 210, 180
31	asaronaldehyde	-	0.5	-	0.8	-	-	M <sup>+</sup> 208, M/z: 96,181,150
32	3,4,5-trimethoxy phenyl-2-	-	6.3	-	0.3	-	1.9	M <sup>+</sup> 224, M/z: 181, 224, 148
33	heptadecanol	-	-	-	1.5	-	1.2	M <sup>+</sup> 256, M/z: 55, 83, 97
34	Citronellol	-	0.9	-	-	-	-	M <sup>+</sup> 156, M/z: 71, 68, 41
35	methyl palmitate	-	-	-	1.8	-	-	M <sup>+</sup> 270, M/z: 74, 87, 143
36	heptadecanol-1	-	1.5	-	1.2	-	-	M <sup>+</sup> 208, M/z: 83, 57
37	phytol	-	-	-	10.6	-	2.3	M <sup>+</sup> 296, M/z: 71,57,123
38	2,13-octadecadien-1-ol	-	-	-	5.0	-	-	M <sup>+</sup> 266, M/z: 55, 41, 67
39	9,12,15-octadecatrienoic acid,	-	-	-	1.3	-	-	M <sup>+</sup> 292, M/z: 79, 67, 41
40	2,4,6 trimethoxy benzonitrile	-	-	-	0.2	-	-	M <sup>+</sup> 193, M/z: m/e: 193, 164, 71
41	leinoleic acid	14.6	-	6.4	-	18.9	-	M <sup>+</sup> 280, M/z: 67, 81, 95
42	Carinol	-	-	-	1.2	-	-	M <sup>+</sup> 378, M/z: 137, 240, 175
43	Pentacosane	0.03	-	-	-	0.2	-	M <sup>+</sup> 352, M/z: 57, 71, 43
44	cyclohexyl palmitate	-	-	-	-	0.2	-	M <sup>+</sup> 338, M/z: 82, 257, 55
45	1,3,12-nonadeca triene	-	-	-	-	1.2	-	M <sup>+</sup> 262, M/z: 67, 41, 82
46	ergost-5-en-3-ol	0.5	-	-	-	0.5	-	M <sup>+</sup> 400, M/z: 107, 96, 145
47	stigmast-4-en-3 one	0.3	0.9	1.0	0.7	1.3	0.5	M <sup>+</sup> 412, M/z: 124, 43, 229
48	vitamin e	0.2	-	0.1	-	0.2	-	M <sup>+</sup> 403, M/z: 165, 430, 164

49	stigmast-5-en-3 ol (3 beta)	3.4	-	3.7	1.3	2.0	-	M <sup>+</sup> 414, M/z: 43, 55, 107
50	4z- decenal	0.1	-	-	-	-	-	M <sup>+</sup> 154, M/z: m/e: 84, 55, 41
51	6-dodecenol	T	-	-	-	-	-	M <sup>+</sup> 184, M/z: 55, 67, 41
52	α- curcumene		0.6	-	0.4	-	-	M <sup>+</sup> 202, M/z: 119, 132, 105
53	Aristolene	0.4	-	-	-	-	-	M <sup>+</sup> 204, M/z: 161, 91
54	β copaene	0.1	-	0.1	-	-	-	M <sup>+</sup> 204, M/z: 161, 105, 91
55	6-epishyobunone	1.2	0.8	-	-	-	2.0	M <sup>+</sup> 220, M/z: 150, 41, 69
56	α- cadinene	-	-	0.2	-	-	-	M <sup>+</sup> 204, M/z: 105,161,204
57	cis methyl isoeugenol	-	-	2	-	-	-	M <sup>+</sup> 178, M/z: 178,142,200
58	α- calacorene	-	-	0.1	-	-	-	M <sup>+</sup> 172, M/z: 157,142,200
59	α- asarone	0.1	4.2	4	3.6	0.2	4.6	M <sup>+</sup> 208, M/z: 208,193,165
60	pentadecanoic acid	-	-	0.2	-	-	-	M <sup>+</sup> 242, M/z: 74, 87, 143
61	methyl linolenate	-	-	0.1	-	-	-	M <sup>+</sup> 292, M/z: 79, 67, 95
62	Tricosane	-	-	0.4	-	-	-	M <sup>+</sup> 324, M/z: 57, 43, 71
63	Stigmasterol	-	-	2.0	-	-	-	M <sup>+</sup> 412, M/z: 55, 83, 69
	Total	90.07	82.00	81.12	80.70	90.75	82.10	-

ACPGLME = *Acorus calamus* Pithoragaeh leaves methanolic extract, ACPNLME = *Acorus calamus* Pantnagar leaves methanolic extract, ACBTLME = *Acorus calamus* Bhimtal leaves methanolic extract, ACPGRME = *Acorus calamus* Pithoragarh rhizomes methanolic extract, ACPNRME = *Acorus calamus* Pantnagar rhizomes methanolic extract, ACBTRME = *Acorus calamus* Bhimtal rhizomes methanolic extract, M<sup>+</sup>= molecular mass, M/z= major fragments, t= trace <.05.

### 3.2. Antioxidant activity

All the extracts exhibited significant DPPH radical scavenging activity as a function of concentration. The radical scavenging potential of the extracts in terms of their IC<sub>50</sub> values revealed the order ACPNRME (108.99 µg/mL) > ACBTRME (206.78 µg/mL) > ACPGLME (298.30µg/mL) > ACBTLME (521.40 µg/mL) > ACPNLME (799.39 µg/mL) > ACPGRME (919.48 µg/mL) compared to the standards catechin (IC<sub>50</sub>= 64.07µg/mL) > BHT (IC<sub>50</sub>= 101.88µg/mL). In the radical form DPPH is purplish/pink in colour and absorb at 517 nm. It has been observed that the radical

scavenger in the DPPH solution donates electrons and reduce the colour of solution and convert it to a stable diamagnetic molecule (Matthaus, 2002). In present investigation the observations were with the total agreement of the above fact. Thus it can be inferred that the phytochemicals in the extracts stabilized the DPPH molecule by donating electrons. It has also been reported that the phenols posses radical scavenging activity and proportionally correlate (Devi and Ganjewala, 2011). In our study the phenols positive correlated between antioxidant activity of the extracts (Table 2 and 3).

**Table 2:** Antioxidant potentials in terms of IC<sub>50</sub> values for different methanolic extracts from *A. calamus* along with their total phenolic content.

Sample/ standard	DPPH scavenging activity/ IC <sub>50</sub> (µg/mL)	Reducing power ability /RP <sub>50</sub> (µg/mL)	Metal chelating ability/ IC <sub>50</sub> (µg/mL)	Total phenols (mg/g GAE)
ACPGLME	298.30±1.25 <sup>d</sup>	177.44±4.80 <sup>c</sup>	270.62±3.52 <sup>b</sup>	3.26±.052 <sup>c</sup>
ACPNLME	799.39±5.75 <sup>f</sup>	105.45±1.34 <sup>a</sup>	202.75±0.19 <sup>a</sup>	2.40±.058 <sup>a</sup>
ACBTLME	521.40±0.82 <sup>e</sup>	267.57±0.34 <sup>e</sup>	314.62±0.06 <sup>bc</sup>	2.45±.051 <sup>b</sup>
ACPGRME	919.48±6.89 <sup>g</sup>	190.50±2.28 <sup>d</sup>	331.30±1.27 <sup>c</sup>	4.10±.055 <sup>e</sup>
ACPNRME	108.99±0.70 <sup>b</sup>	148.91±1.42 <sup>b</sup>	387.78±8.21 <sup>d</sup>	3.41±.052 <sup>d</sup>
ACBTRME	206.78±0.23 <sup>c</sup>	263.21±0.35 <sup>e</sup>	776.23±0.81 <sup>e</sup>	4.80±.054 <sup>f</sup>
BHT	101.88±0.70 <sup>b</sup>	102.23±1.06 <sup>a</sup>	-	-
Catechin	64.077±0.40 <sup>a</sup>	-	-	-
EDTA	-	-	194.06±0.11 <sup>a</sup>	-
Citric acid	-	-	207.56±0.46 <sup>a</sup>	-

ACPGLME = *Acorus calamus* Pithoragaeh leaves methanolic extract, ACPNLME = *Acorus calamus* Pantnagar leaves methanolic extract, ACBTLME = *Acorus calamus* Bhimtal leaves methanolic extract, ACPGRME = *Acorus calamus* Pithoragarh rhizomes methanolic extract, ACPNRME = *Acorus calamus* Pantnagar rhizomes methanolic extract, ACBTRME = *Acorus calamus* Bhimtal rhizomes methanolic extract.- =

Not applicable, Values are means of three replicates  $\pm$  SD, Within a column, mean values followed by the same letter are not significantly different according to Tukey's test ( $p < 0.05$ ), GAE= gallic acid equivalent

It has been reported that antioxidant potential of certain compounds is related to their reducing power and this serves as an important indicator of antioxidant activity. In this assay, ferric ions ( $\text{Fe}^{3+}$ ) were reduced to ferrous ions ( $\text{Fe}^{2+}$ ) with change in colour from yellow to bluish green. The intensity of colour depends on the reducing potential of the compounds present in the reaction medium, consequently, related to the antioxidant activity (Siddhuraju *et al.*, 2002). In present study the  $\text{Fe}^{3+}$  to  $\text{Fe}^{2+}$  reducing activity was determined in terms of their  $\text{RP}_{50}$  values which exhibited the order of ACPNLME (105.45  $\mu\text{g/mL}$ ) > ACPNRME (148.91  $\mu\text{g/mL}$ ) > ACPGLME (177.44  $\mu\text{g/mL}$ ) > ACPGRME (190.50  $\mu\text{g/mL}$ ) > ACBTRME (263.21  $\mu\text{g/mL}$ ) > ACBTLME (267.57  $\mu\text{g/mL}$ ) compared to standard BHT ( $\text{RP}_{50} = 102.23 \mu\text{g/mL}$ ).

Similarly in  $\text{Fe}^{2+}$  metal chelating activity the equilibrium between  $\text{Fe}^{2+}$  ion and ferrozine complex get disturbed in presence of competitor complexing

agent and indicated by decrease in colour intensity thus shows its antioxidant activity (Sethi *et al.*, 2016). In present study the various extracts under investigation revealed varying degree of binding capacity for  $\text{Fe}^{2+}$  ion which expressed their ability as a per oxidation inhibitor. ACPNLME and ACPGLME were found to have significantly higher  $\text{Fe}^{2+}$  chelating abilities followed by other extracts. The  $\text{IC}_{50}$  values for antioxidant potentiality of extracts was obtained as ACPNLME (202.75  $\mu\text{g/mL}$ ) > ACPGLME (270.62  $\mu\text{g/mL}$ ) > ACBTLME (314.62  $\mu\text{g/mL}$ ) > ACPGRME (331.30  $\mu\text{g/mL}$ ) > ACPNRME (387.78  $\mu\text{g/mL}$ ) > ACBTRME (776.23  $\mu\text{g/mL}$ ) compared to standards EDTA and citric acid which exhibited the  $\text{IC}_{50}$  values 194.06  $\mu\text{g/mL}$  and 207.56  $\mu\text{g/mL}$  respectively. The  $\text{IC}_{50}$  values of all the extracts, assayed for their *in-vitro* antioxidant activity by various methods along with their phenolic contents have been reported in table 2, while the correlation of phenols with  $\text{IC}_{50}$  has been recorded in table 3.

**Table 3:** Correlation of total phenols with  $\text{IC}_{50}$  values.

Phenols	Total phenols	DPPH Scavenging activity		Reducing power ability		Metal chelating ability	
		rhizomes	leaves	rhizomes	leaves	rhizomes	leaves
Total phenols	1.0	-0.901	-0.859	-0.998	-0.014	-0.317	0.172

The quantitative estimation of total phenols in all the extracts were observed in the range of 2.40 to 4.80  $\mu\text{g/mL}$ . The highest phenolic content was observed in ACBTRME followed by ACPGRME, ACPNRME, ACPGLME, ACBTLME and ACPNLME. It has already been reported that the phenols in various medicinal herbs play important role as antioxidants (Skerget *et al.*, 2005). From present study it can be inferred that the phenols might be possibly responsible for antioxidant activity in extracts. The total phenols were correlated with  $\text{IC}_{50}/\text{RP}_{50}$  values of different methods used for the determination of antioxidant assay. The results obtained showed negative correlation with DPPH radical scavenging, metal chelating ability and reducing power ability at  $\alpha = 0.01$  or 0.05 (level of significance) except the metal chelating ability of methanolic extract of *A. calamus* leaves (Table 3).

Various biological properties, such as anti-inflammatory, antimicrobial, enzyme inhibition, anti-allergic, and antioxidant activities have been reported to be associated with phenolics and flavanoids (Raina *et al.*, 2003; Shahidi *et al.*, 1992). It has been reported that the hydroxyl group ( $-\text{OH}$ ) in aromatic ring, mediates redox reaction, and is capable of scavenging of free radicals (Raina *et al.*, 2003). Based on the results obtained in present study and traditional use of *A. calamus* as herbal folk medicine it can be concluded that this plant may be a good source of phytochemicals like shynone, linoleic acid, phytol and phenyl propanoids etc. The potential of antioxidant activity in all the extracts and their phenolic content supports its use as safe natural antioxidants and food preservative over the harmful synthetic ones like BHT. However further clinical trials will be needed for its

commercialization and value addition. The study will be also helpful to generate the scientific data base for the researchers so that the herb can be exploited from nature judiciously.

#### 4. Conclusion

Based on above observations observed in present investigation it can be inferred that the herb may be a good source for natural food preservative to prevent the oxidative deteriorative activity of food materials beside generation of database for its scientific and judicious *in-situ* exploitation.

#### 5. Acknowledgements

Authors are thankful to Jawahar Lal University, New Delhi, India for providing GC/MS facility. DST, New Delhi is thankfully acknowledged for financial assistance.

#### 6. References

- Adams, R.P. 2007. Identification of Essential Oil Components by Gas Chromatography/Mass Spectroscopy, 4th Edn. Allured Publishing Co., Carol Stream Illinois.
- Adfa, M., Livandri, F., Meita, N.P., Manaf, S., Ninomiya, M., Gustian, I., Putranto, A.M.H., Supriati, R. and Koketsu, M. 2015. Termiticidal activity of *Acorus calamus* Linn. rhizomes and its main constituents against *Coptotermes curvignathus* Holmgren. *Journal of Asia-Pacific Entomology*, 18: 47-50.
- Avadhani, M.N., Selvaraj, C.I., Rajasekharan, P.E., Rao, V.K., Munirajappa, H. and Tharachand, C. 2016. Genetic diversity analysis and chemical profiling of Indian *Acorus calamus* accessions from South and North-East India. *Indian Journal of Biotechnology*, 4: 560-567.
- Barua, C.C., Sen, S., Das, A.S., Talukdar, A., Hazarika, N.J., Barua, A.G., Barua, A.M. and Barua, I.A. 2014. Comparative study of the in vitro antioxidant property of different extracts of *Acorus calamus* Linn. *Journal of Natural Product and Plant Resources*, 4: 8-18
- Burits, M. and Bucar, F. 2000. Antioxidant activity of *Nigella sativa* essential oil. *Phytotherapy Research*, 14: 323-328.
- Chen, H.P., Yang, K., Zheng, L.S., You, C.X., Cai, Q. and Wang, C.F. 2015. Repellent and insecticidal activities of shyobunone and isoshyobunone derived from the essential oil of *Acorus calamus* rhizomes. *Pharmacognosy Magazine*, 11: 675-681.
- Darmstadt, G.L., Mao-Qiang, M., Chi, E., Saha, S.K., Ziboh, V.A., Black, R.E., Santosham, M. and Elias, P.M. 2002. Impact of tropical oils on the skin barrier: possible implications for neonatal health in developing countries. *Acta Paediatrica*, 91: 546-554.
- Devi, S.A. and Ganjewala, D. 2011. Antioxidant activities of methanolic extracts of sweet-flag (*Acorus calamus*) leaves and rhizomes. *Journal of herbs, spices & medicinal plants*, 17: 1-11.
- Diezel, W., Schulz, E., Shanks, M., and Heise, H. 1993. Plants oils: Topical application and anti-inflammatory effects (Croton oil test). *Dermatologische Monatsschrift*, 179: 173-173.
- Dong, W., Yang, D. and Lu, R. 2010. Chemical constituents from the rhizome of *Acorus calamus* L. *PlantMed*, 76: 454-457.
- Hossain, M.S., Zaman, S., Haque, A.B.M.H., Bhuiyan, M.P.I., Khondkar, P. and Islam, M.R. 2008. Chemical and pesticidal studies on *Acorus calamus* rhizomes. *Journal of Applied Sciences Research*, 4: 1261-1266.
- Jeena, G.S., Punatha, H., Prakash, O., Chandra, M. and Kushwaha, K.P.S. 2016. Study on in vitro antioxidant potential of some cultivated *Pleurotus* species (Oyster mushroom). *Indian Journal of Natural Products and Resources*, 5: 56-61.
- Joshi, N., Prakash, O. and Pant, A.K. 2012. Essential oil composition and in vitro antibacterial activity of rhizome essential oil and  $\beta$ -asarone from *Acorus calamus* L. collected from lower Himalayan Region of Uttarakhand. *Journal of Essential Oil Bearing Plants*, 15: 32-37.
- Kumar, R., Prakash, O., Pant, A.K., Hore, S.K., Chanotiya, C.S. and Mathela, C.S. 2009. Compositional variations and anthelmintic activity of essential oils from rhizomes of different wild populations of *Acorus calamus* L. and its major component, beta-asarone. *Natural Product Communication*, 4: 275-278.
- Kunwar, G., Prakash, O., Chandra, M. and Pant, A.K. 2013. Chemical composition, antifungal and antioxidant activities of *Perilla frutescens* (L.) syn. *P. ocimoides* L. collected from different regions of

- Indian Himalaya. *Asian Journal of Traditional Medicine*, 8:88-98.
- Letawe, C., Boone, M. and Pierard, G. E. 1998. Digital image analysis of the effect of topically applied linoleic acid on acne microcomedones. *Clinical and Experimental Dermatology*, 23: 56-58.
- Lewis, H.W. and Elvin-Lewis, F.P.M. Medical Botany, Plants Affecting Human Health. 2003. Second Edition, 1.
- Matthaus, B. 2002. Antioxidant activity of extracts obtained from residues of different oilseeds. *Journal of Agricultural and Food Chemistry*, 50: 3444-3452.
- Mehrotra, S., Mishra, K.P., Maurya, R., Srimal, R.C., Yadav, V.S., Pandey, R. and Singh, V.K. 2003. Anticellular and immunosuppressive properties of ethanolic extract of *Acorus calamus* rhizome. *International Immunopharmacology*, 3: 53-61.
- Nanda, B.L. 2014. Determination of phytochemicals and antioxidant activity of *Acorus calamus* rhizome. *Journal of Drug Delivery and Therapeutics*, 4: 117-121.
- Pandy, V., Jose, N., Subhash, H. 2009. CNS activity of methanol and acetone extracts of *Acorus calamus* leaves in mice. *Journal of Pharmacology and Toxicology*, 4:79-86.
- Phongpaichit, S., Pujenjob, N., Rukachaisirikul, V. and Ongsakul, M. 2005. Antimicrobial activities of the crude methanol extract of *Acorus calamus* Linn. *Songklanakarin Journal of Science and Technology*, 27: 517-523.
- Raina, V.K., Srivastava, S.K. and Syamasunder, K.V. 2003. Essential oil composition of *Acorus calamus* L. from the lower region of the Himalayas. *Flavour and Fragrance Journal*, 18: 18-20.
- Sethi, S., Prakash, O., Punetha, H. and Pant, A.K. 2016. Antioxidant assay and antifungal activity of essential oil and various extracts from *Alpinia allughas* Roscoe: A potent Zingiberaceous Herb. *Journal of Essential Oil Bearing Plants*, 19: 358-367.
- Shahidi, F., Janitha, P.K. and Wanasudara, P.D. 1992. Phenolic antioxidants. *Critical Reviews in Food Science & Nutrition*; 32: 67-103.
- Shetty, K., Curtis, O.F., Levin, R.E., Witkowsky, R. and Ang, W. 1995. Prevention of vitrification associated with *in vitro* shoot culture of *Oregano* (*Origanum vulgare*) by *Pseudomonas* spp. *Journal of Plant Physiology*, 147: 447-451.
- Siddhuraju, P., Mohan, P.S. and Becker, K. 2002. Studies on the antioxidant activity of Indian Laburnum (*Cassia fistula* L.): a preliminary assessment of crude extracts from stem bark, leaves, flowers and fruit pulp. *Food Chemistry*, 79: 61-67.
- Skerget, M., Kotnik, P., Hadolin, M., Hras, A.R., Simoncic, M. and Knez, Z. 2005. Phenols, proanthocyanidins, flavones and flavonols in some plant materials and their antioxidant activities. *Food Chemistry*, 89: 191-198.